

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrINVOKAMET®

Canagliflozin (as anhydrous canagliflozin) and metformin hydrochloride tablets
Tablets, 50 mg/500 mg, 50 mg/1000 mg, 150 mg/500 mg and 150 mg/1000 mg, Oral

Combinations of oral blood glucose lowering drugs

Janssen Inc.
19 Green Belt Drive
Toronto, Ontario
M3C 1L9

www.janssen.com/canada

All trademarks used under license.

© 2024 Janssen Inc.

Control Number: 280097-01

Date of Initial Authorization:
June 1, 2016

Date of Revision:
June 13, 2024

RECENT MAJOR LABEL CHANGES

2 CONTRAINDICATIONS	06/2023
4 DOSAGE AND ADMINISTRATION, 4.1 Dosing Considerations	03/2024
4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment, Renal Impairment	06/2023
7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism	03/2024
7 WARNINGS AND PRECAUTIONS, Peri-Operative Considerations	03/2024
7 WARNINGS AND PRECAUTIONS, Reproductive Health: Female and Male Potential	03/2024
7 WARNINGS AND PRECAUTIONS, 7.1.1 Pregnant Women	03/2024

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics.....	4
1.2 Geriatrics	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	6
4 DOSAGE AND ADMINISTRATION	7
4.1 Dosing Considerations	7
4.2 Recommended Dose and Dosage Adjustment	7
4.4 Administration.....	8
4.5 Missed Dose.....	8
5 OVERDOSAGE	9
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	10
7 WARNINGS AND PRECAUTIONS	11
7.1 Special Populations	20
7.1.1 Pregnant Women	20
7.1.2 Breast-feeding	20
7.1.3 Pediatrics	21

7.1.4	Geriatrics	21
8	ADVERSE REACTIONS.....	21
8.1	Adverse Reaction Overview	21
8.2	Clinical Trial Adverse Reactions.....	23
8.3	Less Common Clinical Trial Adverse Reactions	28
8.4	Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data	35
8.5	Post-Market Adverse Reactions	39
9	DRUG INTERACTIONS.....	39
9.2	Drug Interactions Overview	39
9.3	Drug-Behavioural Interactions.....	40
9.4	Drug-Drug Interactions	40
9.5	Drug-Food Interactions.....	44
9.6	Drug-Herb Interactions	44
9.7	Drug-Laboratory Test Interactions.....	44
10	CLINICAL PHARMACOLOGY	45
10.1	Mechanism of Action	45
10.2	Pharmacodynamics	46
10.3	Pharmacokinetics	47
11	STORAGE, STABILITY AND DISPOSAL.....	51
12	SPECIAL HANDLING INSTRUCTIONS.....	51
PART II: SCIENTIFIC INFORMATION.....		52
13	PHARMACEUTICAL INFORMATION.....	52
14	CLINICAL TRIALS	53
14.1	Clinical Trials by Indication.....	53
	Type 2 Diabetes Mellitus.....	53
15	MICROBIOLOGY.....	73
16	NON-CLINICAL TOXICOLOGY	73
PATIENT MEDICATION INFORMATION		77

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Note: for additional information on canagliflozin and metformin hydrochloride, consult the individual Product Monographs.

INVOKAMET® (canagliflozin/metformin hydrochloride) is indicated to improve glycemic control as an adjunct to diet and exercise in adult patients with type 2 diabetes mellitus inadequately controlled on:

- metformin
- a sulfonylurea in combination with metformin
- pioglitazone in combination with metformin
- insulin in combination with metformin

Or in patients already being treated and achieving glycemic control with:

- metformin and canagliflozin as separate tablets
- a sulfonylurea in combination with metformin and canagliflozin as separate tablets
- pioglitazone in combination with metformin and canagliflozin as separate tablets
- insulin in combination with metformin and canagliflozin as separate tablets

1.1 Pediatrics

The safety and efficacy of INVOKAMET in pediatric patients under 18 years of age have not been established; therefore, Health Canada has not authorized an indication for pediatric use.

1.2 Geriatrics

Patients 65 years and older had a higher incidence of adverse reactions related to reduced intravascular volume with canagliflozin, including hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration. Reactions were more common in patients over 75 years of age and with the 300 mg daily dose (see [7.1.4 Geriatrics](#) and [4.2 Recommended Dose and Dosage Adjustment](#)). Smaller reductions in HbA1c with canagliflozin relative to placebo were seen in patients 65 years and older, compared to younger patients (see [7.1 Special Populations](#)). Treatment with INVOKAMET can reduce renal function. Metformin is eliminated by the kidney, and the risk of serious adverse reactions to the drug is greater in patients with impaired renal function. INVOKAMET should be used with caution as age increases. The dosage of INVOKAMET should be adjusted based on renal function. Regular assessment of renal function is necessary (see [7 WARNINGS AND PRECAUTIONS](#), [4 DOSAGE AND ADMINISTRATION](#) and [10.3 Pharmacokinetics](#)).

2 CONTRAINDICATIONS

- Known hypersensitivity to canagliflozin or metformin or to any ingredient in the formulation or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).
- In patients with severe renal impairment [estimated glomerular filtration rate (eGFR) <30 mL/min/1.73 m²], end-stage renal disease, in patients on dialysis or when renal function is not known (see [7 WARNINGS AND PRECAUTIONS, Renal](#)).

- Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma, history of ketoacidosis with or without coma, history of lactic acidosis, irrespective of precipitating factors (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#) and [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#)).
- Unstable and/or insulin-dependent (Type I) diabetes mellitus.
- Excessive alcohol intake, acute or chronic.
- Severe hepatic dysfunction, since severe hepatic dysfunction has been associated with some cases of lactic acidosis, INVOKAMET should not be used in patients with clinical or laboratory evidence of hepatic disease (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#)).
- Cardiovascular collapse and in disease states associated with hypoxemia such as cardiorespiratory insufficiency, which are often associated with hyperlactacidemia.
- Stress conditions, such as severe infections, trauma or surgery and the recovery phase thereafter.
- Severe dehydration or shock.
- During pregnancy and breastfeeding (see [7.1.1 Pregnant Women](#) and [7.1.2 Breast-feeding](#))
- Period around administration of iodinated contrast materials, because the use of such products may result in acute alteration of renal function (see [7 WARNINGS AND PRECAUTIONS, Renal](#)).

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

Lactic Acidosis

- Lactic acidosis is a rare, but serious, metabolic complication that can occur due to metformin accumulation during treatment with INVOKAMET (canagliflozin/metformin hydrochloride) (see [7 WARNINGS AND PRECAUTIONS, Lactic Acidosis](#)).
- Patients should be cautioned against excessive alcohol intake, either acute or chronic, when taking INVOKAMET since alcohol intake potentiates the effect of metformin on lactate metabolism (see [7 WARNINGS AND PRECAUTIONS, Lactic Acidosis](#)).

Diabetic Ketoacidosis

- Clinical trial and post-market cases of diabetic ketoacidosis (DKA), a serious life-threatening condition requiring urgent hospitalization, have been reported in patients with type 2 diabetes mellitus (T2DM) treated with canagliflozin, or other sodium-glucose co-transporter 2 (SGLT2) inhibitors. Fatal cases of DKA have been reported in patients taking canagliflozin. A number of cases have been atypical with blood glucose values below 13.9 mmol/L (250 mg/dL) (see [8 ADVERSE REACTIONS](#)).
- The risk of DKA must be considered in the event of non-specific symptoms such as difficulty breathing, nausea, vomiting, abdominal pain, confusion, anorexia, excessive thirst and unusual fatigue or sleepiness. If these symptoms occur, regardless of blood glucose level, INVOKAMET treatment should be immediately **discontinued, and patients should be assessed for DKA immediately**.
- INVOKAMET should not be used for the treatment of DKA or in patients with a history of DKA.
- INVOKAMET is not indicated, and should not be used, in patients with type 1 diabetes.
- See [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#).

Lower Limb Amputation

- An increased risk of lower limb amputations associated with canagliflozin use versus placebo was observed in CANVAS (5.9 vs 2.8 events per 1000 patient-years) and CANVAS-R (7.5 vs 4.2 events per 1000 patient-years), two large, randomized, placebo-controlled trials in patients with type 2 diabetes who had established cardiovascular disease (CVD) or were at risk for CVD.
- Amputations of the toe and midfoot were most frequent; however, amputations involving the leg were also observed. Some patients had multiple amputations, some involving both limbs.
- Before initiating INVOKAMET, consider factors that may increase the risk of amputation, such as a history of prior amputation, peripheral vascular disease, neuropathy, and diabetic foot ulcers.
- Monitor patients receiving INVOKAMET for infection, new pain or tenderness, sores or ulcers involving the lower limbs, and discontinue INVOKAMET if these complications occur.
- See [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

- Renal function should be assessed before initiating INVOKAMET and periodically thereafter (see [7 WARNINGS AND PRECAUTIONS, Renal](#)). In patients with volume depletion not previously treated with canagliflozin, normalize volume status before initiating INVOKAMET (see [7 WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests](#)).
- **Concomitant Use with Insulin or an Insulin Secretagogue (e.g., Sulfonylurea):** When INVOKAMET is used as add-on therapy with insulin or an insulin secretagogue (e.g., sulfonylurea), a lower dose of insulin or the insulin secretagogue may be considered to reduce the risk of hypoglycemia (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#) and [8 ADVERSE REACTIONS](#)).
- **Temporary Interruption for Surgery:** INVOKAMET treatment should be interrupted for a minimum of 3 days, when possible, prior to major surgery or procedures associated with prolonged fasting. Monitor for DKA in the post-operative period. Ensure risk factors for ketoacidosis are resolved and that the patient is clinically stable and has resumed oral intake before considering INVOKAMET treatment re-initiation (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#)).
- INVOKAMET is not recommended for use in patients on loop diuretics (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#) and [8 ADVERSE REACTIONS, Description of Selected Adverse Reaction](#)).

4.2 Recommended Dose and Dosage Adjustment

The dosage of INVOKAMET should be individualized on the basis of both effectiveness and tolerability, while not exceeding the maximum recommended dose of 150 mg canagliflozin/1000 mg metformin hydrochloride twice daily.

INVOKAMET dose escalation should be gradual to reduce the gastrointestinal side effects associated with metformin use.

- In patients on metformin (alone or in combination with a sulfonylurea, pioglitazone, or insulin), switch to INVOKAMET containing canagliflozin 50 mg with a similar total daily dose of metformin;
- In patients already treated with canagliflozin and metformin (alone or in combination with a sulfonylurea, pioglitazone, or insulin), switch to INVOKAMET containing the same total daily doses of each component.

For patients who are tolerating INVOKAMET containing 50 mg canagliflozin taken twice daily who have an eGFR ≥ 60 mL/min/1.73 m², who need tighter glycemic control and who have a low risk of adverse reactions associated with reduced intravascular volume, the dose can be increased to INVOKAMET containing 150 mg canagliflozin taken twice daily (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

Concomitant Use with UDP-Glucuronosyl Transferase (UGT) Enzyme Inducers: If an inducer of UGTs and drug transport systems (e.g., rifampin, phenytoin, barbiturates,

phenobarbital, ritonavir, carbamazepine, efavirenz, St John's wort [*Hypericum perforatum*]) is co-administered with INVOKAMET, monitor HbA1c and consider increasing the dose to canagliflozin 150 mg twice daily in patients currently tolerating 50 mg canagliflozin twice daily who have an eGFR \geq 60 mL/min/1.73 m² or CrCl \geq 60 mL/min and require additional glycemic control. Consider another antihyperglycemic agent in patients with CrCl < 60 mL/min.

Pediatrics (< 18 years of age)

The safety and efficacy of INVOKAMET have not been established in pediatric patients; therefore, Health Canada has not authorized an indication for pediatric use.

Geriatrics (\geq 65 years of age)

Due to the potential for decreased renal function in elderly subjects regular assessment of renal function is necessary (see [7 WARNINGS AND PRECAUTIONS, Renal](#)).

Renal function and risk of volume depletion should be taken into account (see [7 WARNINGS AND PRECAUTIONS](#) and [8 ADVERSE REACTIONS](#)). For those patients who are tolerating canagliflozin 50 mg twice daily and who need tighter glycemic control, the dose can be increased to canagliflozin 150 mg twice daily.

Renal Impairment

No dose adjustment is needed in patients with mild renal impairment (eGFR of 60 mL/min/1.73 m² to < 90 mL/min/1.73 m² or greater).

In patients with eGFR 30 mL/min/1.73 m² to < 60 mL/min/1.73 m² more frequent renal function monitoring is recommended. Metformin daily dose may be reduced with the decline of renal function.

Canagliflozin dose should be limited to 100 mg daily (two 50 mg tablets) in patients with eGFR < 60 mL/min/1.73 m².

In patients with eGFR 30 mL/min/1.73 m² to < 45 mL/min/1.73 m² the maximum daily dose of metformin is 1000 mg.

INVOKAMET is contraindicated and should be discontinued in patients with eGFR < 30 mL/min/1.73 m² (see [2 CONTRAINDICATIONS](#)).

Hepatic Impairment

INVOKAMET is contraindicated in patients with clinical or laboratory evidence of hepatic disease (see [2 CONTRAINDICATIONS](#)). Metformin use in patients with impaired hepatic function has been associated with some cases of lactic acidosis (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#)).

4.4 Administration

INVOKAMET should be taken orally twice a day with meals to reduce the risk of gastrointestinal side effects associated with metformin use. Tablets are to be swallowed whole.

4.5 Missed Dose

If a dose of INVOKAMET is missed, it should be taken as soon as the patient remembers unless

it is almost time for the next dose in which case patients should skip the missed dose and take the medicine at the next regularly scheduled time.

5 OVERDOSAGE

Canagliflozin

Single doses up to 1600 mg of canagliflozin in healthy subjects and canagliflozin 300 mg twice daily for 12 weeks in patients with type 2 diabetes were generally well tolerated.

In the event of an overdose, contact the Poison Control Centre. It is also reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring, and institute supportive treatment as dictated by the patient's clinical status. Canagliflozin was negligibly removed during a 4-hour hemodialysis session. Canagliflozin is not expected to be dialyzable by peritoneal dialysis.

Metformin hydrochloride

Available information concerning treatment of a massive overdose of metformin hydrochloride is very limited. It would be expected that adverse reactions of a more intense character including epigastric discomfort, nausea and vomiting followed by diarrhea, drowsiness, weakness, dizziness, malaise and headache might be seen. Should those symptoms persist, lactic acidosis should be excluded. The drug should be discontinued and proper supportive therapy instituted.

Overdose of metformin hydrochloride has occurred, including ingestion of amounts greater than 50 grams. Hypoglycemia was reported in approximately 10% of cases, but no causal association with metformin hydrochloride has been established. Lactic acidosis has been reported in approximately 32% of metformin overdose cases (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#)). Metformin is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions. Therefore, hemodialysis may be useful for removal of accumulated drug from patients in whom metformin overdose is suspected.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1: Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Tablets 50 mg/500 mg, 50 mg/1000 mg, 150 mg/500 mg, 150 mg/1000 mg	Each tablet contains the following non-medicinal ingredients: <u>Core Tablet:</u> croscarmellose sodium, hypromellose (2910), magnesium stearate, and microcrystalline cellulose. <u>Film Coat:</u> iron oxide black (150 mg/1000 mg tablets, iron oxide red (50 mg/1000 mg, 150 mg/500 mg and 150/1000 mg tablets), iron oxide yellow (50 mg/1000 mg and 150 mg/500 mg), macrogol (polyethylene glycol), polyvinyl alcohol (partially hydrolyzed), talc, and titanium dioxide.

INVOKAMET is supplied as film-coated, immediate-release tablets for oral administration. Each tablet strength contains canagliflozin drug substance as the hemihydrate equivalent to 50 mg or 150 mg of anhydrous canagliflozin, and metformin hydrochloride in quantities of 500 mg or 1000 mg. Tablet strengths are supplied in bottles of 60, inside cartons.

The available tablet strengths are listed below:

Table 2: INVOKAMET Tablet Strengths and Description

Strength	Description
50 mg + 500 mg	Capsule-shaped, white, film-coated tablets with “CM” on one side and “155” on the other side
50 mg + 1000 mg	Capsule shaped, beige, film-coated tablets with “CM” on one side and “551” on the other side
150 mg + 500 mg	Capsule shaped, yellow, film-coated tablets with “CM” on one side and “215” on the other side
150 mg + 1000 mg	Capsule shaped, purple, film-coated tablets with “CM” on one side and “611” on the other side

7 WARNINGS AND PRECAUTIONS

See [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#).

Cardiovascular

Canagliflozin

- Lower limb amputation

An increased risk of lower limb amputations associated with canagliflozin use versus placebo was observed in CANVAS (5.9 and 2.8 amputations per 1000 patients -years) and CANVAS-R (7.5 vs 4.2 events per 1000 patient-years), two large, randomized, placebo-controlled trials evaluating patients with type 2 diabetes who had either established cardiovascular disease or were at risk for cardiovascular disease. The risk of lower limb amputations was observed at both the 100 mg and 300 mg once daily dosage regimens. The amputation data for CANVAS and CANVAS-R are shown in Table 8 and Table 9, respectively (see [8 ADVERSE REACTIONS, Description of Selected Adverse Reactions](#)).

Amputations of the toe and midfoot (99 out of 140 patients with amputations receiving canagliflozin in the two trials) were the most frequent; however, amputations involving the leg, below and above the knee, were also observed (41 out of 140 patients with amputations receiving canagliflozin in the two trials). Some patients had multiple amputations, some involving both lower limbs. Lower limb infections, gangrene, and diabetic foot ulcers were the most common precipitating medical events leading to the need for an amputation. The risk of amputation was highest in patients with a baseline history of prior amputation, peripheral vascular disease, and neuropathy.

Before initiating INVOKAMET, consider factors in the patient history that may predispose to the need for amputations, such as a history of prior amputation, peripheral vascular disease, neuropathy and diabetic foot ulcers. Counsel patients about the importance of routine preventative foot care and adequate hydration. Monitor patients receiving INVOKAMET for signs and symptoms of infection (including osteomyelitis), new pain or tenderness, sores or ulcers involving the lower limbs, and discontinue INVOKAMET if these complications occur.

- Reduced Intravascular Volume

Due to its mechanism of action, canagliflozin increases urinary glucose excretion (UGE) and induces an osmotic diuresis, which may reduce intravascular volume. Patients most susceptible to adverse reactions related to reduced intravascular volume (e.g., postural dizziness, orthostatic hypotension, hypotension or renal failure) include patients with moderate renal impairment, elderly patients, patients on loop diuretics or medications that interfere with the renin-angiotensin-aldosterone system (e.g., angiotensin-converting-enzyme [ACE] inhibitors, angiotensin receptor blockers [ARBs]), and patients with low systolic blood pressure (see [8 ADVERSE REACTIONS, Description of Selected Adverse Reactions](#), [9 DRUG INTERACTIONS](#) and [4 DOSAGE AND ADMINISTRATION](#)). Before initiating INVOKAMET in patients with one or more of these characteristics, volume status should be assessed and any volume depletion corrected. Caution should also be exercised in other patients for whom a drop in blood pressure could pose a risk, such as patients with known cardiovascular disease. Monitor for signs and symptoms after initiating therapy. Patients should be advised to report symptoms of reduced intravascular volume.

In placebo-controlled clinical studies of canagliflozin, increases in adverse reactions related to reduced intravascular volume were seen more commonly with the 300 mg canagliflozin dose and occurred most frequently in the first three months (see [8 ADVERSE REACTIONS, Description of Selected Adverse Reactions](#)).

INVOKAMET is not recommended for use in patients receiving loop diuretics (see [8 ADVERSE REACTIONS, Description of Selected Adverse Reactions](#) and [4.1 Dosing Considerations](#)) or who are volume depleted.

In case of intercurrent conditions that may lead to volume depletion (such as a gastrointestinal illness), careful monitoring of volume status (e.g., physical examination, blood pressure measurements, laboratory tests including renal function tests), and serum electrolytes is recommended. In the case of volume depletion, temporary interruption of treatment with INVOKAMET may be considered until the condition is corrected, and more frequent glucose monitoring may be considered.

Metformin hydrochloride

- Hypoxic States

Cardiovascular collapse (shock) from whatever cause (e.g., acute congestive heart failure, acute myocardial infarction and other conditions characterized by hypoxemia) have been associated with lactic acidosis and may also cause prerenal azotemia. When such events occur in patients on INVOKAMET therapy, the drug should be promptly discontinued.

Driving and Operating Machinery

The effect of canagliflozin on the ability to drive and use machines has not been examined. However, patients should be alerted to the elevated risk of adverse reactions related to reduced intravascular volume, such as postural dizziness, and to the risk of hypoglycemia when INVOKAMET is used as add-on therapy with insulin or an insulin secretagogue (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#) and [Endocrine and Metabolism, 8 ADVERSE REACTIONS](#) and [4.1 Dosing Considerations](#)).

Endocrine and Metabolism

Canagliflozin

- Diabetic ketoacidosis

Clinical trial and post-market cases of DKA, a serious life-threatening condition requiring urgent hospitalization, have been reported in patients with type 2 diabetes mellitus treated with SGLT2 inhibitors, including canagliflozin. Fatal cases of DKA have been reported in patients taking canagliflozin. In a number of reported cases, the presentation of the condition was atypical with blood glucose values below 13.9 mmol/L (250 mg/dL) (see [8 ADVERSE REACTIONS, Description of Selected Adverse Reactions](#)).

INVOKAMET should not be used in patients with type 1 diabetes. The diagnosis of T2DM should therefore be confirmed before initiating INVOKAMET.

INVOKAMET is not indicated and should not be used for the treatment of DKA or in patients with a history of DKA.

Patients with type 2 diabetes treated with INVOKAMET who present with signs and symptoms consistent with severe metabolic acidosis should be assessed for ketoacidosis regardless of presenting blood glucose levels, as ketoacidosis associated with canagliflozin may be present even if blood glucose levels are less than 13.9 mmol/L (250 mg/dL).

The risk of DKA must be considered in the event of non-specific symptoms such as difficulty breathing, nausea, vomiting, abdominal pain, confusion, anorexia, excessive thirst and unusual fatigue or sleepiness.

If these symptoms occur, regardless of blood glucose level, INVOKAMET treatment should be immediately discontinued, patients should be assessed for diabetic ketoacidosis immediately, and prompt treatment should be instituted.

SGLT2 inhibitors have been shown to increase blood ketones in clinical trial subjects. Conditions that can precipitate DKA while taking INVOKAMET include patients on a very low carbohydrate diet (as the combination may further increase ketone body production), patients with conditions that lead to a restricted food intake or severe dehydration, patients with increased insulin requirements due to an acute medical illness, surgery, or alcohol abuse, patients with a low beta-cell function reserve [e.g., type 2 diabetes patients with low C-peptide or latent autoimmune diabetes in adults (LADA)], pancreatic disorders suggesting insulin deficiency (e.g., type 1 diabetes, history of pancreatitis, or pancreatic surgery), insulin dose reduction (including insulin pump failure), and patients with a history of ketoacidosis. These patients should be monitored closely. Caution should also be taken when reducing the insulin dose in patients requiring insulin (see [4 DOSAGE AND ADMINISTRATION](#)).

Prolonged diabetic ketoacidosis

DKA may be prolonged in some patients. In most post-marketing adverse event reports, ketoacidosis lasted for 3 days or more despite INVOKAMET discontinuation and standard treatment of diabetic ketoacidosis. Based on canagliflozin half-life, glucosuria may persist longer than expected and DKA may be prolonged in some patients. The mechanism of prolonged DKA and glucosuria is unknown. Other factors independent of canagliflozin may be involved in prolonging periods of DKA. In post-marketing adverse event reports, most cases reported prolongation that lasted from 3 to 10 days after discontinuation of INVOKAMET, however, a few cases reported longer prolongation (see [8.5 Post-Market Adverse Reactions](#)).

Treatment interruption considerations

Temporarily discontinue treatment with INVOKAMET in T2DM patients who are hospitalized for major surgical procedures, or will undergo scheduled surgery, and in patients who are hospitalized for serious infections or acute serious medical illnesses. INVOKAMET treatment should be interrupted for a minimum of 3 days, when possible, prior to major surgery or any other procedures associated with prolonged fasting, when, based on the drug half-life, most of INVOKAMET would be expected to be eliminated. Monitoring for DKA is recommended in these patients even if drug treatment has been interrupted or discontinued. Ensure risk factors for ketoacidosis are resolved prior to considering INVOKAMET treatment re-initiation (See [4.1. Dosing Considerations](#) and [7 WARNINGS AND PRECAUTIONS, Peri-operative Conditions](#)).

Educate patients on the signs and symptoms of ketoacidosis and instruct patients to discontinue INVOKAMET and seek medical attention immediately if signs and symptoms occur.

- Hypoglycemia in Add-on Therapy with other Antihyperglycemic Agents

When canagliflozin was used as add-on therapy with insulin or an insulin secretagogue (e.g., sulfonylurea), the incidence of hypoglycemia was increased over that of placebo. Therefore, to lower the risk of hypoglycemia, a dose reduction of insulin or an insulin secretagogue may be considered (see [8 ADVERSE REACTIONS](#) and [4.1 Dosing Considerations](#)).

- Increases in Low-Density Lipoprotein (LDL-C)

Dose-related increases in LDL-C are seen with canagliflozin treatment (see [8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data](#)) LDL-C levels should be monitored.

Metformin hydrochloride

- Hypoglycemia

Hypoglycemia does not occur in patients receiving metformin alone under usual circumstances of use, but could occur when caloric intake is deficient, when strenuous exercise is not compensated by caloric supplementation, or during concomitant use with other glucose lower agents or ethanol.

Elderly, debilitated, or malnourished patients and those with adrenal or pituitary insufficiency or alcohol intoxication are particularly susceptible to hypoglycemic effects. Hypoglycemia may be difficult to recognize in the elderly, and in people who are taking β -adrenergic blocking drugs.

- Lactic Acidosis

Lactic acidosis is a rare, but serious, metabolic complication that occurs due to metformin accumulation during treatment with INVOKAMET. When it occurs, it is fatal in approximately 50% of cases. Lactic acidosis may also occur in association with a number of pathophysiologic conditions, including diabetes mellitus, and whenever there is significant tissue hypoperfusion and hypoxemia. Lactic acidosis is characterized by elevated blood lactate levels (> 5 mmol/L), decreased blood pH, electrolyte disturbances with an increased anion gap, and an increased lactate/pyruvate ratio. When metformin is implicated as the cause of lactic acidosis, metformin plasma levels > 5 μ g/mL are generally found.

The reported incidence of lactic acidosis in patients receiving metformin HCl is very low (approximately 0.03 cases / 1000 patient-years, with approximately 0.015 fatal cases / 1000 patient-years) and occurs primarily in diabetic patients with significant renal insufficiency, including both intrinsic renal disease and renal hypoperfusion, often in the setting of multiple concomitant medical/surgical problems and multiple concomitant medications. Patients with congestive heart failure requiring pharmacologic management, in particular those with unstable or acute congestive heart failure who are at risk of hypoperfusion and hypoxemia, are at increased risk of lactic acidosis. In particular, treatment of the elderly should be accompanied by careful monitoring of renal function. INVOKAMET treatment should not be initiated in patients ≥ 80 years of age, unless measurement of creatinine clearance demonstrates that renal function is not reduced, as the patients are more susceptible to developing lactic acidosis. The risk of lactic acidosis increases with the degree of renal dysfunction and the patient's age. The risk of lactic acidosis may, therefore, be significantly decreased by regular monitoring of renal function in patients taking metformin and by use of the minimum effective dose of metformin. In addition, INVOKAMET should be promptly withheld in the presence of any condition associated with hypoxemia, dehydration or sepsis. Because impaired hepatic function may significantly limit the

ability to clear lactate, INVOKAMET should generally be avoided in patients with clinical or laboratory evidence of hepatic disease. Patients should be cautioned against excessive alcohol intake, either acute or chronic, when taking metformin hydrochloride, since alcohol intake potentiates the effect of metformin hydrochloride on lactate metabolism. In addition, INVOKAMET should be temporarily discontinued prior to any intravascular radiocontrast study and for any surgical procedure.

The onset of lactic acidosis is often subtle, and accompanied only by nonspecific symptoms such as malaise, myalgia, respiratory distress, increasing somnolence and non-specific abdominal distress. There may be associated hypothermia, hypotension and resistance bradyarrhythmias with more marked acidosis. The patient and the patient's physician must be aware of the possible importance of such symptoms and the patient should be instructed to notify the physician immediately if they occur. INVOKAMET should be withdrawn until the situation is clarified. Serum electrolytes, ketones, blood glucose and, if indicated, blood pH, lactate levels and even blood metformin levels may be useful. Once a patient is stabilized on any dose level of metformin, gastrointestinal symptoms, which are common during initiation of therapy, are unlikely to be drug related. Later occurrence of gastrointestinal symptoms could be due to lactic acidosis or other serious disease. In patients taking metformin, levels of fasting venous plasma lactate above the upper limit of normal but <5 mmol/L, do not necessarily indicate impending lactic acidosis and may be explainable by other mechanisms, such as poorly controlled diabetes or obesity, vigorous physical activity or technical problems in sample handling. Lactic acidosis should be suspected in any diabetic patient with metabolic acidosis lacking evidence of ketoacidosis (ketonuria and ketonemia).

Lactic acidosis is a medical emergency that must be treated in a hospital setting. In a patient with lactic acidosis who is taking INVOKAMET, the drug should be discontinued immediately and general supportive measures should be promptly instituted. Because metformin hydrochloride is dialysable (with clearance of up to 170 mL/min under good hemodynamic conditions), prompt hemodialysis is recommended to correct the acidosis and remove the accumulated metformin.

Physicians should instruct their patients to recognize the symptoms which could be signal onset of lactic acidosis. If acidosis of any kind develops, INVOKAMET should be discontinued immediately and the patient should be immediately hospitalized.

- **Change in Clinical Status in Patients with Previously Controlled Type 2 Diabetes**

A patient with type 2 diabetes previously well-controlled on INVOKAMET who develops laboratory abnormalities or clinical illness (especially vague and poorly defined illness) should be evaluated promptly for evidence of ketoacidosis or lactic acidosis. Evaluation should include serum electrolytes and ketones, blood glucose and, if indicated, blood pH, lactate, pyruvate, and metformin levels. If acidosis of either form occurs, INVOKAMET must be stopped immediately and other appropriate corrective measures initiated.

- **Loss of control of blood glucose**

When a patient stabilized on any diabetic regimen is exposed to stress such as fever, trauma, infection, or surgery, a temporary loss of glycemic control may occur. At such times, it may be necessary to withhold INVOKAMET and temporarily administer insulin. INVOKAMET may be reinstated after the acute episode is resolved.

- Vitamin B₁₂ levels

In controlled clinical trials of metformin of 29-week duration, a decrease to subnormal levels of previously normal serum Vitamin B₁₂ levels, without clinical manifestations, was observed in approximately 7% of patients. Such decrease, possibly due to interference with B₁₂ absorption from the B₁₂-intrinsic factor complex, is, however, very rarely associated with anemia and appears to be rapidly reversible with discontinuation of metformin or Vitamin B₁₂ supplementation. Measurement of hematologic parameters on an annual basis is advised in patients on metformin and any apparent abnormalities should be appropriately investigated and managed (see [7 WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests](#) and [8.1 Adverse Reaction Overview](#)). Certain individuals (those with inadequate Vitamin B₁₂ or calcium intake or absorption) appear to be predisposed to developing subnormal Vitamin B₁₂ levels. Measurements of serum Vitamin B₁₂ are advisable at least every one to two years in patients on long term INVOKAMET therapy.

Genitourinary

Canagliflozin

- Genital Mycotic Infections

Canagliflozin increases the risk of genital mycotic infections, consistent with the mechanism of increased urinary glucose. Patients with a history of genital mycotic infections and uncircumcised males were more likely to develop genital mycotic infections (see [8 ADVERSE REACTIONS](#)).

- Urinary tract infections (including urosepsis and pyelonephritis)

Treatment with canagliflozin increases the risk for urinary tract infections (see [8 ADVERSE REACTIONS](#)). There have been post-marketing reports of serious urinary tract infections, including urosepsis and pyelonephritis, requiring hospitalization in patients treated with canagliflozin.

- Fournier's gangrene (necrotizing fasciitis of the perineum)

Post-marketing cases of necrotizing fasciitis of the perineum (Fournier's gangrene), a rare but serious and potentially life-threatening necrotizing infection requiring urgent surgical intervention, have been reported in female and male patients with diabetes mellitus receiving SGLT2 inhibitors, including INVOKAMET. Serious outcomes have included hospitalization, multiple surgeries, and death.

Patients treated with INVOKAMET who present with pain or tenderness, erythema, or swelling in the genital or perineal area, with or without fever, or malaise should be evaluated for necrotizing fasciitis. If suspected, INVOKAMET should be discontinued and prompt treatment should be instituted (including broad-spectrum antibiotics and surgical debridement if necessary).

Hematologic

Canagliflozin

Mean hemoglobin and hematocrit increased in patients administered canagliflozin, as did the frequency of patients with abnormally elevated values for hemoglobin/hematocrit (see [8.4](#)

[Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data](#)). INVOKAMET should be used with caution in patients with an elevated hematocrit.

Hepatic/Biliary/Pancreatic

INVOKAMET is contraindicated in patients with clinical or laboratory evidence of hepatic disease (see [2 CONTRAINDICATIONS](#)). Impaired hepatic function has been associated with some cases of lactic acidosis. INVOKAMET has not been studied in patients with hepatic impairment.

Canagliflozin

Canagliflozin has not been studied in patients with severe hepatic impairment.

Metformin hydrochloride

Metformin use in patients with impaired hepatic function has been associated with some cases of lactic acidosis.

Immune

Serious hypersensitivity reactions, including angioedema and anaphylaxis, have been reported post-market in patients treated with canagliflozin. If a hypersensitivity reaction is suspected, discontinue INVOKAMET, assess for other potential causes and initiate alternative treatment for diabetes (see [8.5 Post-Market Adverse Reactions](#)).

Monitoring and Laboratory Tests

Blood Glucose and HbA1c

Response to INVOKAMET treatment should be monitored by periodic measurements of blood glucose and HbA1c levels. Due to its mechanism of action, patients taking INVOKAMET will test positive for glucose in their urine.

Renal function

Renal function should be assessed prior to initiation of INVOKAMET and regularly thereafter (see [4 DOSAGE AND ADMINISTRATION](#)). INVOKAMET is contraindicated in patients with severe renal impairment [estimated glomerular filtration rate (eGFR) <30 mL/min/1.73 m²], end-stage renal disease, in patients on dialysis or when renal function is not known (see [2 CONTRAINDICATIONS](#) and [4.2 Recommended Dose and Dosage Adjustment, Renal Impairment](#)).

Reduced intravascular volume:

INVOKAMET is not recommended for use in patients who are volume depleted. Before initiating INVOKAMET, assess volume status, particularly in patients at risk (e.g., moderate renal impairment, the elderly, in patients with low systolic blood pressure, or if on a loop diuretic, angiotensin-converting enzyme inhibitor, or angiotensin receptor blocker).

In patients with volume depletion, the condition should be corrected prior to initiation of INVOKAMET (see [4.1 Dosing Considerations](#)).

For patients with risk factors for volume depletion or in case of intercurrent conditions that may lead to volume depletion (such as a gastrointestinal illness), careful monitoring of volume status

(e.g., physical examination, blood pressure measurements, laboratory tests including renal function tests), and serum electrolytes is recommended during treatment with INVOKAMET. Temporary interruption of treatment with INVOKAMET should be considered until volume depletion is corrected.

LDL-cholesterol

LDL-C levels should be measured at baseline and at regular intervals during treatment with INVOKAMET due to dose-dependent increases in LDL-C seen with therapy.

Digoxin levels

In patients taking digoxin and canagliflozin 300 mg once daily for seven days, there was an increase in the total exposure (AUC) and peak drug concentration (C_{max}) of digoxin (20% and 36%, respectively), therefore patients taking INVOKAMET concomitantly with digoxin should be monitored appropriately.

Hematology

Initial and periodic monitoring of hematologic parameters (e.g., hemoglobin/hematocrit and red blood cell indices) should be performed regularly. While megaloblastic anemia has rarely been seen with metformin therapy, if this is suspected, Vitamin B₁₂ deficiency should be excluded.

Musculoskeletal

Canagliflozin

An increased risk of bone fracture, occurring as early as 12 weeks after treatment initiation, was observed in patients using canagliflozin in the CANVAS cardiovascular outcomes trial. Consider factors that contribute to fracture risk prior to initiating canagliflozin (see [8 ADVERSE REACTIONS, Bone fractures](#)).

Peri-Operative Considerations

Temporarily discontinue treatment with INVOKAMET for a minimum of 3 days in T2DM patients who are hospitalized for major surgical procedures, or will undergo scheduled surgery. Ensure risk factors for ketoacidosis are resolved and that the patient is clinically stable and has resumed oral intake before considering INVOKAMET treatment re-initiation. Monitoring for DKA is recommended in these patients (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#) and [4.1 Dosing Considerations](#)).

Renal

Impairment of renal function

Canagliflozin increases serum creatinine and decreases eGFR in a dose dependent fashion. In clinical trials, renal function abnormalities have occurred after initiating canagliflozin. Post-marketing cases of acute kidney injury, including acute renal failure and a decline in eGFR, some requiring hospitalization and dialysis, shortly after initiation of canagliflozin treatment have been reported. Before initiating INVOKAMET, consider factors that may predispose patients to acute kidney injury including hypovolemia, chronic renal insufficiency, congestive heart failure and concomitant medications (diuretics, ACE inhibitors, ARBs, NSAIDs) (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#) and [8 ADVERSE REACTIONS](#)). Consider temporarily discontinuing INVOKAMET in any setting of reduced oral intake (such as acute illness or fasting) or fluid losses (such as gastrointestinal illness or excessive heat exposure); monitor

patients for signs and symptoms of acute kidney injury. If acute kidney injury occurs, discontinue INVOKAMET promptly and institute treatment.

Renal function should be assessed prior to initiation of INVOKAMET and regularly thereafter. More frequent renal function monitoring is recommended in patients whose eGFR decreases to < 60 mL/min/1.73 m² after initiating treatment.

Metformin is excreted by the kidney, and the risk of metformin accumulation and lactic acidosis increases with the degree of impairment of renal function. Thus, patients with serum creatinine levels above the upper limit of the normal range for their age should not receive INVOKAMET. Before initiation of INVOKAMET therapy and every 6 months while on INVOKAMET therapy, renal function should be assessed and verified as being within normal range.

Decreased renal function occurs more commonly in elderly patients and can be asymptomatic. In patients in whom development of renal dysfunction is anticipated, renal function should be assessed more frequently and INVOKAMET discontinued if evidence of renal impairment is present.

Special caution should be exercised in situations where renal function may become impaired, for example when initiating antihypertensive therapy or diuretic therapy and when starting therapy with an NSAID.

Use of Concomitant Medications that May Affect Renal Function or Metformin Disposition

Concomitant medication(s) that may affect renal function or result in a significant hemodynamic change or interfere with the disposition of metformin such as cationic drugs that are eliminated by renal tubular secretion should be used with caution (see [9 DRUG INTERACTIONS](#)).

Administration of Iodinated Contrast Agent

Intravascular contrast studies with iodinated materials can lead to acute alteration of renal function and have been associated with lactic acidosis in patients receiving metformin (see [2 CONTRAINDICATIONS](#)). Therefore, in patients in whom any such study is planned, INVOKAMET should be temporarily discontinued at the time of or prior to the procedure, and withheld for 48 hours subsequent to the procedure and reinstated only after renal function has been re-evaluated and found to be normal.

Use in renal impairment

INVOKAMET is contraindicated in patients with severe renal impairment [estimated glomerular filtration rate (eGFR) < 30 mL/min/1.73 m²], end-stage renal disease, in patients on dialysis or when renal function is not known (see [2 CONTRAINDICATIONS](#) and [4.2 Recommended Dose and Dosage Adjustment, Renal Impairment](#)).

Reproductive Health: Female and Male Potential

See [2 CONTRAINDICATIONS](#) and [7.1.1 Pregnant women](#).

- **Fertility**

No human fertility studies have been conducted. No animal studies have been conducted with the combined products in INVOKAMET to evaluate impairment of fertility. For the effects on fertility of the individual components of INVOKAMET, see [16 NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology](#).

- **Teratogenic Risk**

INVOKAMET is contraindicated during pregnancy (see [2 CONTRAINDICATIONS](#) and [7.1.1 Pregnant Women](#)). There are no adequate and well-controlled studies of INVOKAMET or its components in pregnant women, and therefore, the safety of INVOKAMET in human fetal development has not been established. Based on animal studies, canagliflozin may affect renal development and maturation. In animal studies, adverse renal effects were observed in juvenile rats when canagliflozin was administered during a period of renal development corresponding to the late second and third trimesters of human pregnancy (see [16 NON-CLINICAL TOXICOLOGY, Juvenile Toxicity](#)).

7.1 Special Populations

7.1.1 Pregnant Women

INVOKAMET is contraindicated in pregnancy (see [2 CONTRAINDICATIONS](#)). There are no adequate and well-controlled studies in pregnant women with INVOKAMET or its individual components, therefore, the safety of INVOKAMET in pregnant women is not known.

Canagliflozin

Based on results from rat studies, canagliflozin may affect renal development and maturation. In a juvenile rat study, increased kidney weights, and renal pelvic and tubular dilatation that did not fully reverse within the 1-month recovery period, were evident at greater than or equal to 0.5 times clinical exposure from a 300 mg dose (see [16 NON-CLINICAL TOXICOLOGY, Juvenile Toxicity](#)).

Metformin hydrochloride

Metformin was not teratogenic in rats and rabbits at doses up to 600 mg/kg/day, or about two times the maximum recommended human daily dose on a body surface area basis. Determination of fetal concentrations demonstrated a partial placental barrier to metformin. Because animal reproduction studies are not always predictive of human response, metformin is contraindicated during pregnancy (see [2 CONTRAINDICATIONS](#)).

7.1.2 Breast-feeding

INVOKAMET is contraindicated during nursing (see [2 CONTRAINDICATIONS](#)). No studies in lactating animals have been conducted with INVOKAMET or its individual components.

Canagliflozin

It is not known if canagliflozin is excreted in human milk. Available pharmacodynamic/toxicological data in animals have shown excretion of canagliflozin in the milk of lactating rats reaching levels which are approximately 1.4 times higher than plasma systemic

exposure. Data in juvenile rats directly exposed to canagliflozin showed risk to the developing kidney (renal pelvic and tubular dilatations) during maturation.

Metformin hydrochloride

Studies in lactating rats show that metformin is excreted into milk and reaches levels comparable to those in plasma. Metformin is excreted into human milk.

7.1.3 Pediatrics

Safety and effectiveness of INVOKAMET in pediatric patients under 18 years of age have not been established; therefore, Health Canada has not authorized an indication for pediatric use.

7.1.4 Geriatrics

INVOKAMET treatment is associated with reduced renal function and should be used with caution as age increases because elderly patients are more likely to have decreased renal function (see [7 WARNINGS AND PRECAUTIONS, Renal](#) and [4 DOSAGE AND ADMINISTRATION, Geriatrics](#)).

Canagliflozin

Two thousand thirty-four (2,034) patients 65 years and older, and 345 patients 75 years and older were exposed to canagliflozin in nine clinical studies of canagliflozin (see [14 CLINICAL TRIALS](#)).

Patients 65 years and older had a higher incidence of adverse reactions related to reduced intravascular volume with canagliflozin (such as hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration), particularly with the 300 mg daily dose, compared to younger patients; more prominent increase in the incidence was seen in patients who were 75 years and older (see [4 DOSAGE AND ADMINISTRATION](#) and [8 ADVERSE REACTIONS](#)). Smaller reductions in HbA1c with canagliflozin relative to placebo were seen in older (65 years and older; -0.61% with canagliflozin 100 mg and -0.74% with canagliflozin 300 mg relative to placebo) compared to younger patients (-0.72% with canagliflozin 100 mg and -0.87% with canagliflozin 300 mg relative to placebo).

Metformin hydrochloride

Controlled clinical studies of metformin did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients, although other reported clinical experience has not identified differences in responses between the elderly and younger patients. Metformin is known to be substantially excreted by the kidney and because the risk of serious adverse reactions to the drug is greater in patients with impaired renal function, metformin should only be used in patients with normal renal function (see [2 CONTRAINDICATIONS](#)). Because aging is associated with reduced renal function, INVOKAMET should be used with caution as age increases.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

There have been no clinical studies conducted with INVOKAMET tablets. INVOKAMET tablets demonstrated comparable bioavailability of canagliflozin and metformin with co-administered tablets of canagliflozin and metformin in comparative bioavailability studies (see [10.3](#)

[Pharmacokinetics](#)).

Canagliflozin

The safety of canagliflozin was evaluated in fifteen double-blind, controlled Phase 3 and Phase 4 clinical studies involving 22,645 patients with type 2 diabetes, including 13,278 patients treated with canagliflozin 100 mg and 7,170 patients, treated with canagliflozin 300 mg. Of the 22,645 patients with type 2 diabetes, a total of 10,134 patients were treated in two dedicated cardiovascular outcomes studies, for a mean exposure duration of 149 weeks (223 weeks in CANVAS and 94 weeks in CANVAS-R), and 8,114 patients were treated in 12 double-blind, controlled Phase 3 and Phase 4 clinical studies, for a mean exposure duration of 49 weeks. In a dedicated renal outcomes study, a total of 4,397 patients with type 2 diabetes and diabetic nephropathy had a mean duration of drug exposure of 115 weeks.

The primary assessment of safety and tolerability was conducted in a pooled analysis (N=2313) of four 26-week placebo-controlled clinical studies (monotherapy and add-on therapy with metformin, metformin and sulfonylurea, and metformin and pioglitazone). The most commonly reported adverse reactions during treatment ($\geq 5\%$) were vulvovaginal candidiasis, urinary tract infection (UTI), and polyuria or pollakiuria. Adverse reactions leading to discontinuation of $\geq 0.5\%$ of all canagliflozin-treated patients in these studies were vulvovaginal candidiasis (0.7% of females) and balanitis or balanoposthitis (0.5% of males).

A total of 8 serious adverse drug reactions were reported in the primary placebo-controlled safety population, including 5 reports from patients taking canagliflozin 100 mg daily (2 urticaria, 2 UTI, and 1 nausea), 2 reports from patients taking canagliflozin 300 mg daily (1 UTI, 1 constipation) and 1 report from a patient in the placebo group (reduced intravascular volume). Of these serious adverse reactions, 2 led to discontinuation in the canagliflozin group (UTI and urticaria).

Metformin hydrochloride

Lactic acidosis: very rare ($< 1/10,000$ and isolated reports) (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#), and [5 OVERDOSAGE](#)).

Gastrointestinal Reactions: very common ($> 1/10$). Gastrointestinal symptoms (diarrhea, nausea, vomiting, abdominal bloating, flatulence, and anorexia) are the most common reactions to metformin and are approximately 30% more frequent in patients on metformin monotherapy than in placebo-treated patients, particularly during initiation of metformin therapy. These symptoms are generally transient and resolve spontaneously during continued treatment. Because significant diarrhea and/or vomiting can cause dehydration and prerenal azotemia, metformin should be temporarily discontinued, under such circumstances.

For patients who have been stabilized on metformin, nonspecific gastrointestinal symptoms should not be attributed to therapy unless intercurrent illness or lactic acidosis have been excluded.

Special Senses: common ($\geq 1/100$): taste disturbance, i.e. metallic taste.

Dermatologic Reactions: very rare ($< 1/10,000$ and isolated reports): Reports of skin reactions such as erythema, pruritus, and urticaria are very rare.

Hematologic: During controlled clinical trials of 29 weeks duration, approximately 9% of patients

on metformin monotherapy developed asymptomatic subnormal serum Vitamin B₁₂ levels; serum folic acid levels did not decrease significantly. Five cases of megaloblastic anemia have been reported with metformin administration and no increased incidence of neuropathy has been observed. (See [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#)).

Metformin can commonly reduce vitamin B₁₂ serum levels which may result in clinically significant vitamin B₁₂ deficiency (e.g., megaloblastic anemia). Consideration of such etiology is recommended if a patient presents with megaloblastic anemia. The risk of low vitamin B₁₂ levels increases with increasing metformin dose, treatment duration, and/or in patients with risk factors known to cause vitamin B₁₂ deficiency. Periodic monitoring of vitamin B₁₂ levels is recommended in these patients.

Hepatic: very rare (<1/10,000 and isolated reports). Liver function tests abnormalities or hepatitis resolving upon metformin discontinuation has been documented in isolated reports.

Canagliflozin and Metformin hydrochloride

The incidence and type of adverse reactions in 26-week placebo-controlled metformin add-on studies were similar to the adverse reactions in the four 26-week placebo-controlled clinical studies used for the primary assessment of safety and tolerability. There were no additional adverse reactions identified in the pooling of these three placebo-controlled studies that included metformin relative to the four placebo-controlled studies.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Table 3 to Table 6 include treatment-emergent adverse events (TEAEs) reported in ≥ 2% of canagliflozin-treated patients.

Combination with Metformin (Studies DIA3006 and DIA3009)

Table 3: Adverse events (regardless of causality) reported in ≥ 2% of patients treated with canagliflozin and more frequently than in the placebo groups* in double-blind clinical trials of canagliflozin in add-on combination use with metformin, and compared to sitagliptin or placebo (Study DIA3006, 26 weeks) or to glimepiride (Study DIA3009, 52 weeks)

System Organ Class / Preferred Term	Study DIA3006 (26 weeks)				Study DIA3009 (52 weeks)		
	Placebo + Metformin n=183 n (%)	Canagliflozin 100 mg + Metformin n=368 n (%)	Canagliflozin 300 mg + Metformin N=367 n (%)	Sitagliptin 100 mg + Metformin n=366 n (%)	Canagliflozin 100 mg + Metformin n=483 n (%)	Canagliflozin 300 mg + Metformin n=485 n (%)	Glimepiride + Metformin n=482 n (%)
Gastrointestinal Disorders							
Diarrhea	12 (6.6)	12 (3.3)	18 (4.9)	16 (4.4)	24 (5.0)	33 (6.8)	29 (6.0)
Gastritis	3 (1.6)	3 (0.8)	8 (2.2)	3 (0.8)	2 (0.4)	5 (1.0)	7 (1.5)
Nausea	3 (1.6)	11 (3.0)	8 (2.2)	5 (1.4)	16 (3.3)	25 (5.2)	13 (2.7)

System Organ Class / Preferred Term	Study DIA3006 (26 weeks)				Study DIA3009 (52 weeks)		
	Placebo + Metformin n=183 n (%)	Canagliflozin 100 mg + Metformin n=368 n (%)	Canagliflozin 300 mg + Metformin N=367 n (%)	Sitagliptin 100 mg + Metformin n=366 n (%)	Canagliflozin 100 mg + Metformin n=483 n (%)	Canagliflozin 300 mg + Metformin n=485 n (%)	Glimepiride + Metformin n=482 n (%)
Toothache	2 (1.1)	3 (0.8)	8 (2.2)	4 (1.1)	8 (1.7)	7 (1.4)	6 (1.2)
Vomiting	1 (0.5)	8 (2.2)	1 (0.3)	3 (0.8)	9 (1.9)	7 (1.4)	8 (1.7)
General Disorders and Administration Site Conditions							
Fatigue	2 (1.1)	10 (2.7)	8 (2.2)	1 (0.3)	9 (1.9)	7 (1.4)	10 (2.1)
Pyrexia	3 (1.6)	4 (1.1)	5 (1.4)	3 (0.8)	11 (2.3)	9 (1.9)	7 (1.5)
Thirst	0	2 (0.5)	4 (1.1)	0	8 (1.7)	14 (2.9)	0
Infections and Infestations							
Bronchitis	2 (1.1)	2 (0.5)	5 (1.4)	9 (2.5)	11 (2.3)	9 (1.9)	10 (2.1)
Gastroenteritis	2 (1.1)	3 (0.8)	3 (0.8)	2 (0.5)	3 (0.6)	15 (3.1)	9 (1.9)
Influenza	5 (2.7)	6 (1.6)	4 (1.1)	8 (2.2)	17 (3.5)	17 (3.5)	8 (1.7)
Sinusitis	3 (1.6)	8 (2.2)	2 (0.5)	6 (1.6)	7 (1.4)	13 (2.7)	6 (1.2)
Urinary Tract Infection	4 (2.2)	19 (5.2)	13 (3.5)	12 (3.3)	27 (5.6)	24 (4.9)	18 (3.7)
Vaginal Infection	0	2 (0.5)	3 (0.8)	1 (0.3)	11 (2.3)	7 (1.4)	1 (0.2)
Vulvovaginal Mycotic Infection	0	10 (2.7)	7 (1.9)	1 (0.3)	6 (1.2)	14 (2.9)	4 (0.8)
Musculoskeletal and Connective Tissue Disorders							
Back Pain	6 (3.3)	8 (2.2)	12 (3.3)	4 (1.1)	29 (6.0)	18 (3.7)	20 (4.1)
Musculoskeletal Pain	1 (0.5)	3 (0.8)	6 (1.6)	5 (1.4)	9 (1.9)	10 (2.1)	9 (1.9)
Psychiatric Disorders							
Insomnia	0	3 (0.8)	0	1 (0.3)	7 (1.4)	10 (2.1)	6 (1.2)
Renal and Urinary Disorders							
Pollakiuria	1 (0.5)	21 (5.7)	10 (2.7)	2 (0.5)	12 (2.5)	12 (2.5)	1 (0.2)
Reproductive System and Breast Disorders							
Balanoposthitis	1 (0.5)	2 (0.5)	1 (0.3)	0	4 (0.8)	13 (2.7)	2 (0.4)
Vulvovaginal Pruritus	0	4 (1.1)	5 (1.4)	1 (0.3)	6 (1.2)	20 (4.1)	1 (0.2)

*In either study

Combination with a Metformin and a Sulfonylurea (Studies DIA3002 and DIA3015)

Table 4: Adverse events (regardless of causality) reported in $\geq 2\%$ of patients treated with canagliflozin and more frequently than in the placebo groups* in double-blind clinical trials of canagliflozin in add-on combination use with metformin and a sulfonylurea, and compared to placebo (Study DIA3002, 26 weeks) or sitagliptin (Study DIA3015, 52 weeks)

System Organ Class / Preferred Term	Study DIA3002 (26 weeks)			Study DIA3015 (52 weeks)	
	Placebo+ Metformin + Sulfonylurea n=156 n (%)	Canagliflozin 100 mg + Metformin + Sulfonylurea n=157 n (%)	Canagliflozin 300 mg + Metformin + Sulfonylurea N=156 n (%)	Canagliflozin 300 mg + Metformin + Sulfonylurea n=377 n (%)	Sitagliptin 100 mg+ Metformin + Sulfonylurea n=378 n (%)
Ear and Labyrinth Disorders					
Vertigo	1 (0.6)	1 (0.6)	1 (0.6)	14 (3.7)	11 (2.9)
Gastrointestinal Disorders					
Abdominal Pain	1 (0.6)	2 (1.3)	1 (0.6)	8 (2.1)	6 (1.6)
Abdominal Pain Upper	2 (1.3)	1 (0.6)	1 (0.6)	10 (2.7)	2 (0.5)
Constipation	0	4 (2.5)	5 (3.2)	9 (2.4)	3 (0.8)
Diarrhea	5 (3.2)	5 (3.2)	10 (6.4)	17 (4.5)	26 (6.9)
Nausea	1 (0.6)	2 (1.3)	4 (2.6)	9 (2.4)	11 (2.9)
Infections and Infestations					
Bronchitis	3 (1.9)	4 (2.5)	3 (1.9)	1 (0.3)	11 (2.9)
Influenza	7 (4.5)	2 (1.3)	3 (1.9)	22 (5.8)	15 (4.0)
Nasopharyngitis	4 (2.6)	6 (3.8)	8 (5.1)	33 (8.8)	38 (10.1)
Sinusitis	3 (1.9)	4 (2.5)	2 (1.3)	8 (2.1)	8 (2.1)
Tooth Abscess	0	4 (2.5)	1 (0.6)	0	2 (0.5)
Upper Respiratory Tract Infection	10 (6.4)	17 (10.8)	6 (3.8)	33 (8.8)	21 (5.6)
Urinary Tract Infection	8 (5.1)	9 (5.7)	8 (5.1)	15 (4.0)	19 (5.0)
Vulvovaginal Mycotic Infection	2 (1.3)	8 (5.1)	8 (5.1)	12 (3.2)	5 (1.3)
Metabolism and Nutrition Disorders					
Decreased Appetite	1 (0.6)	0	4 (2.6)	4 (1.1)	5 (1.3)
Hypoglycemia	6 (3.8)	11 (7.0)	9 (5.8)	66 (17.5)	75 (19.8)
Musculoskeletal and Connective Tissue Disorders					
Arthralgia	4 (2.6)	7 (4.5)	7 (4.5)	17 (4.5)	8 (2.1)
Back Pain	4 (2.6)	2 (1.3)	5 (3.2)	8 (2.1)	15 (4.0)
Musculoskeletal Pain	1 (0.6)	0	3 (1.9)	8 (2.1)	6 (1.6)
Nervous System Disorders					
Headache	4 (2.6)	5 (3.2)	2 (1.3)	29 (7.7)	27 (7.1)
Renal and Urinary Disorders					
Pollakiuria	1 (0.6)	4 (2.5)	3 (1.9)	6 (1.6)	5 (1.3)
Reproductive System and Breast Disorders					
Vulvovaginal Pruritus	0	1 (0.6)	3 (1.9)	15 (4.0)	1 (0.3)

*In either study

Combination with Metformin and Pioglitazone (Study DIA3012)

Table 5: Adverse events (regardless of causality) reported in $\geq 2\%$ of patients treated with canagliflozin and more frequently than in the placebo group in a 26-week double-blind clinical trial of canagliflozin in add-on combination use with metformin and pioglitazone, and compared to placebo (Study DIA3012)

SYSTEM ORGAN CLASS / Preferred Term	Placebo + Metformin+ Pioglitazone n=115 n (%)	Canagliflozin 100 mg + Metformin + Pioglitazone n=113 n (%)	Canagliflozin 300 mg + Metformin + Pioglitazone n=114 n (%)
Gastrointestinal Disorders			
Gastritis	2 (1.7)	4 (3.5)	0
General Disorders And Administration Site Conditions			
Fatigue	2 (1.7)	1 (0.9)	4 (3.5)
Edema Peripheral	2 (1.7)	2 (1.8)	4 (3.5)
Thirst	0	5 (4.4)	4 (3.5)
Infections And Infestations			
Nasopharyngitis	6 (5.2)	6 (5.3)	11 (9.6)
Sinusitis	2 (1.7)	1 (0.9)	3 (2.6)
Upper Respiratory Tract Infection	7 (6.1)	9 (8.0)	5 (4.4)
Vulvovaginal Candidiasis	0	1 (0.9)	3 (2.6)
Vulvovaginal Mycotic Infection	0	3 (2.7)	6 (5.3)
Investigations			
Weight Decreased	1 (0.9)	1 (0.9)	3 (2.6)
Metabolism And Nutrition Disorders			
Hypoglycemia	2 (1.7)	1 (0.9)	6 (5.3)
Musculoskeletal And Connective Tissue Disorders			
Arthralgia	2 (1.7)	1 (0.9)	6 (5.3)
Back Pain	3 (2.6)	8 (7.1)	5 (4.4)
Pain in Extremity	1 (0.9)	4 (3.5)	3 (2.6)
Nervous System Disorders			
Dizziness	1 (0.9)	4 (3.5)	3 (2.6)
Headache	4 (3.5)	3 (2.7)	5 (4.4)
Renal And Urinary Disorders			
Pollakiuria	1 (0.9)	5 (4.4)	7 (6.1)
Reproductive System And Breast Disorders			
Balanitis	0	3 (2.7)	0
Respiratory, Thoracic and Mediastinal Disorders			
Oropharyngeal Pain	2 (1.7)	3 (2.7)	0
Vascular Disorders			
Hypotension	3 (2.6)	3 (2.7)	0

Combination with Insulin and Metformin (Study DIA3008 Insulin Substudy)

Table 6: Adverse events (regardless of causality) reported in $\geq 2\%$ of patients treated with canagliflozin and more frequently than in the placebo group in a 18-week double-blind clinical trial of canagliflozin in add-on combination use with insulin and metformin, and compared to placebo (Study DIA3008 - Insulin Substudy)

System Organ Class / Preferred Term	Placebo + Insulin + Metformin n=244 n (%)	Canagliflozin 100 mg + Insulin + Metformin n=241 n (%)	Canagliflozin 300 mg + Insulin + Metformin n=246 n (%)
Gastrointestinal disorders			
Constipation	2 (0.8)	1 (0.4)	8 (3.3)
Diarrhea	7 (2.9)	4 (1.7)	14 (5.7)
Dyspepsia	0	2 (0.8)	5 (2.0)
Nausea	5 (2.0)	5 (2.1)	8 (3.3)
General disorders and administration site conditions			
Fatigue	4 (1.6)	6 (2.5)	8 (3.3)
Thirst	0	2 (0.8)	10 (4.1)
Infections and infestations			
Bronchitis	5 (2.0)	7 (2.9)	3 (1.2)
Nasopharyngitis	22 (9.0)	22 (9.1)	13 (5.3)
Urinary tract infection	4 (1.6)	3 (1.2)	10 (4.1)
Vulvovaginal mycotic infection	2 (0.8)	4 (1.7)	5 (2.0)
Metabolism and nutrition disorders			
Hypoglycemia	21 (8.6)	23 (9.5)	23 (9.3)
Musculoskeletal and connective tissue disorders			
Arthralgia	3 (1.2)	8 (3.3)	4 (1.6)
Back pain	5 (2.0)	3 (1.2)	13 (5.3)
Pain in extremity	4 (1.6)	7 (2.9)	6 (2.4)
Nervous system disorders			
Dizziness	0	1 (0.4)	6 (2.4)
Headache	7 (2.9)	8 (3.3)	7 (2.8)
Renal and urinary disorders			
Pollakiuria	1 (0.4)	7 (2.9)	18 (7.3)
Reproductive system and breast disorders			
Balanitis	1 (0.4)	7 (2.9)	9 (3.7)
Vascular disorders			
Hypertension	3 (1.2)	8 (3.3)	1 (0.4)

8.3 Less Common Clinical Trial Adverse Reactions

Below is a list of less common (< 2%)¹ clinical trial adverse reactions.

Metabolism and nutrition disorders: dehydration²

Nervous system disorders: dizziness postural², syncope²

Skin and subcutaneous tissue disorders: rash³, urticaria

Vascular disorders: hypotension², orthostatic hypotension²

Description of Selected Adverse Reactions

Reduced intravascular volume: In the pooled analysis of the four 26-week, placebo-controlled studies, the incidence of all adverse reactions related to reduced intravascular volume (e.g., postural dizziness, orthostatic hypotension, hypotension, dehydration, and syncope) was 1.2% for canagliflozin 100 mg, 1.3% for canagliflozin 300 mg, and 1.1% for placebo. The incidence of these adverse reactions with canagliflozin treatment in the two active-controlled studies was similar to comparators.

In one of the dedicated long-term cardiovascular studies (CANVAS), where patients were generally older with a higher prevalence of comorbidities, the incidence rate of adverse reactions related to reduced intravascular volume were 2.34 with canagliflozin 100 mg, 2.87 with canagliflozin 300 mg, and 1.85 with placebo, events per 100 patient-years of exposure.

In the long-term renal outcomes trial, the incidence of hypotension was 2.8% in the canagliflozin 100 mg group and 1.5% in the placebo group.

To assess risk factors for these adverse reactions, a larger pooled analysis (N=12,441) of patients from 13 controlled Phase 3 and Phase 4 studies including both doses of canagliflozin was conducted. In this pooled analysis, patients on loop diuretics, patients with moderate renal impairment (eGFR 30 to < 60 mL/min/1.73 m²), and patients ≥ 75 years of age had higher incidences of these reactions. For patients on loop diuretics, the incidence rates were 4.98 on canagliflozin 100 mg and 5.67 on canagliflozin 300 mg compared to 4.15 events per 100 patient-years of exposure in the control group. For patients with a baseline eGFR 30 to < 60 mL/min/1.73 m², the incidence rates were 5.24 on canagliflozin 100 mg and 5.35 on canagliflozin 300 mg compared to 3.11 events per 100 patient-years of exposure in the control group. In patients ≥ 75 years of age, the incidence rates were 5.27 on canagliflozin 100 mg and 6.08 on canagliflozin 300 mg compared to 2.41 events per 100 patient-years of exposure in the control group (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#), [4 DOSAGE AND ADMINISTRATION](#) and [10 CLINICAL PHARMACOLOGY, Special Populations and Conditions](#)).

¹ Adverse drug reactions (ADRs) were identified based on a comprehensive assessment of biological plausibility, mechanism of action, dose dependence in incidence rate, time of onset, seriousness and consistency of findings across four, 26-week placebo-controlled Phase 3 clinical studies. Additional supportive safety analyses were conducted on a large pooled dataset from eight active- and placebo-controlled Phase 3 clinical studies.

² Related to reduced intravascular volume (see **Adverse reactions related to reduced intravascular volume**).

³ Rash includes the terms: rash erythematous, rash generalized, rash macular, rash maculopapular, rash papular, rash pruritic, rash pustular, and rash vesicular

Diabetic ketoacidosis: Cases of DKA, a serious life-threatening condition requiring urgent hospitalization, have been reported in patients with type 2 diabetes mellitus treated with SGLT2 inhibitors, including canagliflozin. In the on-treatment analysis of the CANVAS/CANVAS-R integrated dataset, the adjusted incidence rates of adjudicated diabetic ketoacidosis were 0.08 (0.2%, 14/5,790) and 0.01 (<0.1%, 1/4,344) per 100 subject-years, for the combined canagliflozin and the placebo groups, respectively. Fatal cases of DKA have been reported in patients treated with canagliflozin. INVOKAMET should not be used in patients with type 1 diabetes. In a number of reported cases, the presentation of the condition was atypical with blood glucose values below 13.9 mmol/L (250 mg/dL) (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#)).

In a long-term renal outcomes study in patients with type 2 diabetes and diabetic nephropathy, on-treatment incidence rates of adjudicated events of DKA were 0.22 (0.5%, 11/2,200) and 0.02 (<0.1%, 1/2,197) per 100 patient-years with canagliflozin 100 mg and placebo, respectively; of the 12 patients with DKA, 7 (6 on canagliflozin 100 mg and 1 on placebo) had an eGFR before treatment of 30 to < 45 mL/min/1.73 m². Cases of DKA in the canagliflozin group occurred in the setting of an intercurrent illness requiring hospitalization (8 of 11 subjects), or with low beta cell function reserve (3 of 11 subjects).

Hypoglycemia: In individual clinical trials (see [14 CLINICAL TRIALS](#)), episodes of hypoglycemia occurred at a higher rate when canagliflozin was co-administered with insulin or sulfonylurea (see Table 7; [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#) and [4 DOSAGE AND ADMINISTRATION](#)).

Table 7: Incidence of Hypoglycemia¹ in Controlled Clinical Studies

Monotherapy (26 weeks)	Placebo (N=192)	Canagliflozin 100 mg (N=195)	Canagliflozin 300 mg (N=197)
Overall [N (%)]	5 (2.6)	7 (3.6)	6 (3.0)
In Combination with Metformin (26 weeks)	Placebo + Metformin (N=183)	Canagliflozin 100 mg + Metformin (N=368)	Canagliflozin 300 mg + Metformin (N=367)
Overall [N (%)]	3 (1.6)	16 (4.3)	17 (4.6)
Severe [N (%)] ²	0 (0)	1 (0.3)	1 (0.3)
In Combination with Metformin (52 weeks)	Glimepiride + Metformin (N=482)	Canagliflozin 100 mg + Metformin (N=483)	Canagliflozin 300 mg + Metformin (N=485)
Overall [N (%)]	165 (34.2)	27 (5.6)	24 (4.9)
Severe [N (%)] ²	15 (3.1)	2 (0.4)	3 (0.6)
In Combination with Metformin + Sulfonylurea (26 weeks)	Placebo + Metformin + Sulfonylurea (N=156)	Canagliflozin 100 mg + Metformin + Sulfonylurea (N=157)	Canagliflozin 300 mg + Metformin + Sulfonylurea (N=156)
Overall [N (%)]	24 (15.4)	43 (27.4)	47 (30.1)
Severe [N (%)] ²	1 (0.6)	1 (0.6)	0
In Combination with Metformin + Sulfonylurea (52 weeks)	Sitagliptin + Metformin + Sulfonylurea (N=378)		Canagliflozin 300 mg + Metformin + Sulfonylurea (N=377)
Overall [N (%)]	154 (40.7)		163 (43.2)
Severe [N (%)] ²	13 (3.4)		15 (4.0)
In Combination with Metformin + Pioglitazone (26 weeks)	Placebo + Metformin + Pioglitazone (N=115)	Canagliflozin 100 mg + Metformin + Pioglitazone (N=113)	Canagliflozin 300 mg + Metformin + Pioglitazone (N=114)
Overall [N (%)]	3 (2.6)	3 (2.7)	6 (5.3)
In Combination with Insulin (18 weeks)	Placebo + Insulin (N=565)	Canagliflozin 100 mg + Insulin (N=566)	Canagliflozin 300 mg + Insulin (N=587)
Overall [N (%)]	208 (36.8)	279 (49.3)	285 (48.6)
Severe [N (%)] ²	14 (2.5)	10 (1.8)	16 (2.7)
In Combination with Insulin and Metformin (18 weeks)³	Placebo + Metformin + Insulin (N=244)	Canagliflozin 100 mg + Metformin + Insulin (N=241)	Canagliflozin 300 mg + Metformin + Insulin (N=246)
Overall [N (%)]	101(41.1)	107 (44.4)	113 (45.9)
Severe [N (%)] ²	9 (3.7)	4 (1.7)	4 (1.6)

¹ Number of patients experiencing at least one event of hypoglycemia based on either biochemically documented episodes (any glucose value ≤ 3.89 mmol/L) or severe hypoglycemic events in the intent-to-treat population.

² Severe episodes of hypoglycemia were defined as those where the patient: required the assistance of another person to recover; lost consciousness; or experienced a seizure (regardless of whether biochemical documentation of a low glucose value was obtained).

³ Subgroup of patients (N=731) from insulin substudy on canagliflozin in combination with metformin and insulin

Twice Daily Dosing

The incidence of hypoglycemia in a Phase 2 clinical study with twice daily dosing (canagliflozin 50 mg or 150 mg twice daily in combination with metformin) was reported in 4.3% and 3.2% of patients treated with canagliflozin 50 mg and 150 mg twice daily, respectively, compared to 3.2% in placebo-treated patients. There were no cases of severe hypoglycemia reported in the canagliflozin or placebo groups.

Genital mycotic infections: Vulvovaginal candidiasis (including vulvovaginitis and vulvovaginal mycotic infection) was reported in 10.4% and 11.4% of female patients treated with canagliflozin 100 mg and canagliflozin 300 mg, respectively, compared to 3.2% in placebo-treated female patients. Most reports of vulvovaginal candidiasis occurred during the first four months of treatment with canagliflozin. Among female patients taking canagliflozin, 2.3% experienced more than one infection. Overall, 0.7% of all female patients discontinued canagliflozin due to vulvovaginal candidiasis (see [7 WARNINGS AND PRECAUTIONS, Genitourinary](#)).

Candidal balanitis or balanoposthitis was reported in 4.2% and 3.7% of male patients treated with canagliflozin 100 mg and canagliflozin 300 mg, respectively, compared to 0.6% in placebo-treated male patients. Among male patients taking canagliflozin, 0.9% had more than one infection. Overall, 0.5% of male patients discontinued canagliflozin due to candidal balanitis or balanoposthitis. In uncircumcised males in a pooled analysis of 10 controlled studies, the incidence rate of phimosis was 0.56 events per 100 patient-years of exposure in patients treated with canagliflozin and 0.05 events per 100 patient-years in patients treated with comparator. In this pooled analysis, the incidence rate of circumcision was 0.38 events per 100 patient-years of exposure in male patients treated with canagliflozin compared to 0.10 events per 100 patient-years in male patients treated with comparator (see [7 WARNINGS AND PRECAUTIONS, Genitourinary](#)).

In the CANVAS integrated dataset, the adjusted-incidence rates of any male mycotic genital infection were 3.17 and 0.96 per 100 patient-years in the combined canagliflozin and placebo groups, respectively.

Urinary tract infections: Urinary tract infections were more frequently reported for canagliflozin 100 mg and 300 mg (5.9% versus 4.3%, respectively) compared to 4.0% with placebo. Most infections were mild to moderate with no increase in the occurrence of serious adverse events. Subjects responded to standard treatments while continuing canagliflozin treatment. The incidence of recurrent infections was not increased with canagliflozin.

Fournier's gangrene (necrotizing fasciitis of the perineum): Fournier's gangrene was identified as a SGLT2i class adverse reaction based on spontaneous event reporting. These events had not been previously identified as ADRs because there were very few subjects in the canagliflozin Phase 3 and Phase 4 clinical development program (including the CANVAS and CREDENCE programs) with adverse events of Fournier's gangrene (incidences were <0.1% in the canagliflozin and comparator groups). All 4 events of Fournier's gangrene (2 subjects treated with canagliflozin and 2 subjects treated with comparator) in the canagliflozin Phase 3 and Phase 4 clinical development program were serious.

Falls: In the pool of all Phase 3 studies, the incidence rate of AEs coded as related to a fall was 7.3, 8.0, and 11.8 per 1000 patient years of exposure to comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

Bone fractures: In cardiovascular study (CANVAS) of 4,327 patients with established or at least two risk factors for cardiovascular disease, the incidence rates of all adjudicated bone fracture were 1.59, 1.79, and 1.09 per 100 patient-years of follow up to canagliflozin 100 mg, canagliflozin 300 mg, and placebo, respectively, with the fracture imbalance initially occurring within the first 26 weeks of therapy.

In two other long-term studies (CANVAS-R and CREDENCE) and in studies conducted in the general diabetes population, no difference in fracture risk was observed with canagliflozin relative to control. In CANVAS-R, of 5,807 patients with established or at least two risk factors for cardiovascular disease, the incidence rates of all adjudicated bone fracture were 1.14 and 1.32 events per 100 patient-years of follow up to canagliflozin and placebo, respectively.

In a long-term renal outcomes study (CREDENCE) of 4,397 patients with type 2 diabetes and diabetic nephropathy, the incidence rates of all adjudicated bone fracture were 1.18 and 1.21 events per 100 patient-years of follow-up for canagliflozin 100 mg and placebo, respectively. In other type 2 diabetes studies with canagliflozin, which enrolled a general diabetes population of 7,729 patients and where bone fractures were adjudicated, the incidence rates of all adjudicated bone fracture were 1.18 and 1.08 events per 100 patient-years of follow up to canagliflozin and control, respectively.

Decreases in Bone Mineral Density: Bone mineral density (BMD) was measured by dual-energy X-ray absorptiometry in a clinical trial of 714 older adults (mean age 64 years). At 2 years, patients randomized to canagliflozin 100 mg and canagliflozin 300 mg had placebo-corrected declines in BMD at the total hip of 0.9% and 1.2%, respectively, and at the lumbar spine of 0.3% and 0.7%, respectively. Placebo-adjusted BMD declines were 0.1% at the femoral neck for both canagliflozin doses and 0.4% at the distal forearm for patients randomized to canagliflozin 300 mg. The placebo-adjusted change at the distal forearm for patients randomized to canagliflozin 100 mg was 0%.

Photosensitivity: In the CANVAS outcome trials integrated dataset, the adjusted-incidence rates of photosensitivity adverse events were 1.03 (0.3%, 19/5790) and 0.26 (0.1%, 3/4344) events per 1,000 subject-years in the combined canagliflozin and the placebo groups, respectively. In a dataset from 12 other phase 3 or 4 trials (excluding the CANVAS outcome trials) that enrolled a diabetic population of 8114 patients, an imbalance in phototoxicity adverse events was not seen with canagliflozin relative to control.

Skin ulcers and peripheral ischemia: In the pool of 8 clinical studies with 78 weeks of mean duration of exposure, skin ulcers occurred in 0.7%, 1.1%, and 1.5% of patients and peripheral ischemia occurred in 0.1%, 0.4%, and 0.2% of patients receiving comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. An imbalance in these events generally were seen within the first 24 weeks of treatment and occurred in patients with known or at high risk for atherosclerotic disease, longer duration of diabetes, presence of diabetic complications, and diuretic use. In the on-treatment analysis set of the CREDENCE renal outcomes trial, there was a higher incidence rate of adverse events of diabetic foot reported in the canagliflozin group compared with the placebo group: 8.47 (43 subjects) and 4.89 (24 subjects) per 1,000 subject-years, respectively.

Renal Cell Carcinoma: In the CANVAS outcome trials integrated dataset, the adjusted-incidence rates of any renal cell carcinoma adverse event were 0.62 (0.2%, 14/5790) and 0.21 (0.1%, 3/4344) per 1,000 subject-years in the canagliflozin and the placebo groups, respectively. Whether this numerical imbalance is related to canagliflozin treatment is unknown.

Lower limb amputation: An increased risk of lower limb amputations associated with canagliflozin use was observed in CANVAS (5.9 vs 2.8 events per 1000 patient years) and CANVAS-R (7.5 vs 4.2 events per 1000 patient-years), two large, randomized, placebo-controlled trials evaluating patients with type 2 diabetes who had either established cardiovascular disease or were at risk for cardiovascular disease. The imbalance occurred as early as the first 26 weeks of therapy. Patients in CANVAS and CANVAS-R were followed for an average of 5.7 and 2.1 years, respectively. The amputation data for CANVAS and CANVAS-R are shown in Table 8 and Table 9, respectively. See [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#).

Table 8: CANVAS Amputations

	Placebo (N=1441)	Canagliflozin 100 mg (N=1445)	Canagliflozin 300 mg (N=1441)	Canagliflozin pooled (N=2886)
Patients with an amputation, n (%)	22 (1.5)	50 (3.5)	45 (3.1)	95 (3.3)
Total amputations	33	83	79	162
Amputation incidence rate (per 1000 patient-years)	2.8	6.2	5.5	5.9
Hazard ratio (95% CI)		2.24 (1.36, 3.69)	2.01 (1.20, 3.34)	2.12 (1.34, 3.38)

Note: Incidence is based on the number of patients with at least one amputation, and not the total number of amputation events. A patient's follow-up is calculated from Day 1 to the first amputation event date. Some patients had more than one amputation

Table 9: CANVAS-R Amputations

	Placebo (N=2903)	Canagliflozin 100 mg (with up-titration to 300 mg) (N=2904)
Patients with an amputation, n (%)	25 (0.9)	45 (1.5)
Total amputations	36	59
Amputation incidence rate (per 1000 patient-years)	4.2	7.5
Hazard Ratio (95% CI)	--	1.80 (1.10, 2.93)

Note: Incidence is based on the number of patients with at least one amputation, and not the total number of amputation events. A patient's follow-up is calculated from Day 1 to the first amputation event date. Some patients had more than one amputation.

In a data pool of patients from 12 other phase 3 or 4 trials (excluding CANVAS program) that enrolled a diabetic population of 8114 patients, the majority of which were without cardiovascular disease, no difference in lower limb amputation risk was observed on canagliflozin relative to control.

The risk of lower limb amputations associated with the use of canagliflozin 100 mg relative to placebo was 12.3 vs 11.2 events per 1000 patient-years, respectively in CREDENCE, a long-term renal outcomes study of 4,397 patients with type 2 diabetes and diabetic nephropathy, with a mean follow-up duration of 136 weeks (see Table 10 and [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

Table 10: Lower limb amputations CREDENCE (On-study analysis)

	Placebo (N=2197)	Canagliflozin 100 mg (N=2200)
Patients with an amputation, n (%)	63 (2.9)	70 (3.2)
Total amputations	96	87
Amputation incidence rate (per 1000 patient-years)	11.2	12.3
Hazard Ratio (95% CI)	--	1.11 (0.79, 1.56)

Note: Incidence is based on the number of patients with at least one amputation, and not the total number of amputation events. A patient's follow-up is calculated from Day 1 to the first amputation event date. Some patients had more than one amputation.

Elderly Patients: Compared to younger patients, patients 65 years and older had a higher incidence of adverse reactions related to reduced intravascular volume with canagliflozin, including hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration. In particular, in patients ≥ 75 years of age, adverse reactions related to reduced intravascular volume occurred with incidence rates of 5.27, 6.08, and 2.41 events per 100 patient-years of exposure for canagliflozin 100 mg, canagliflozin 300 mg, and the control group, respectively. Decreases in eGFR (-3.41 and -4.67 mL/min/1.73 m²) were reported with canagliflozin 100 mg and 300 mg, respectively, compared to the control group (-4.15 mL/min/1.73 m²) (see [7 WARNINGS AND PRECAUTIONS](#) and [4.2 Recommended Dose and Dosage Adjustment](#)).

Patients with Type 2 Diabetes Mellitus and an eGFR 45 to < 60 mL/min/1.73 m²: In a pooled analysis of patients (N=722) with a baseline eGFR 45 to < 60 mL/min/1.73 m², the incidence rates of adverse reactions related to reduced intravascular volume were 4.61 for canagliflozin 100 mg and 4.37 for canagliflozin 300 mg relative to 3.00 events per 100 patient-years of exposure for placebo (see [7 WARNINGS AND PRECAUTIONS](#) and [4 DOSAGE AND ADMINISTRATION](#)). Serum creatinine levels increased from baseline to end of treatment by 5.92 and 6.98 $\mu\text{mol/L}$ for canagliflozin 100 mg and 300 mg, respectively, relative to 7.03 $\mu\text{mol/L}$ with placebo. Blood urea nitrogen (BUN) levels increased by from baseline to end of treatment by 0.92 and 0.77 $\mu\text{mol/L}$ for canagliflozin 100 mg and 300 mg, respectively, relative to 0.57 $\mu\text{mol/L}$ with placebo. The incidence rates of decreases in eGFR (< 80 mL/min/1.73 m² and > 30% decrease from baseline) at any time during treatment were 5.17, 6.62, and 5.82 events per 100 patient-years of exposure for canagliflozin 100 mg, canagliflozin 300 mg, and placebo, respectively. At the last post-baseline value, incidence rates for such decreases were 2.52 for patients treated with canagliflozin 100 mg, 1.91 for patients treated with canagliflozin 300 mg, and 3.20 events per 100 patient-years of exposure for placebo (see [7 WARNINGS AND PRECAUTIONS](#)).

The incidences of elevated serum potassium (> 5.4 mEq/L and 15% above baseline) at any post-baseline value were 4.11 for canagliflozin 100 mg, 4.33 for canagliflozin 300 mg, and 3.8 events per 100 patient-years of exposure for placebo. Rare, more severe elevations were seen in patients with moderate renal impairment who had prior elevated potassium

concentrations and/or who were on multiple medications that reduce potassium excretion, such as potassium-sparing diuretics and angiotensin-converting-enzyme (ACE) inhibitors.

Serum phosphate changes from baseline to end of treatment were 0.00 and 0.02 mmol/L for canagliflozin 100 mg and 300 mg, respectively, compared to 0.00 mmol/L for placebo. The incidence rates of elevated serum phosphate (> 1.65 mmol/L and 25% above baseline) at any post-baseline value were 0.93 for canagliflozin 100 mg, 1.15 for canagliflozin 300 mg, and 0.71 events per 100 patient-years of exposure for placebo.

Patients with Type 2 Diabetes Mellitus and an eGFR 30 to < 60 mL/min/1.73 m² Treated for Diabetic Nephropathy: In a long-term renal outcomes study in patients with type 2 diabetes and diabetic nephropathy, the incidence rate for renal-related adverse events was lower in the canagliflozin 100 mg group compared with the placebo group (7.23 and 10.55 per 100 patient-years in canagliflozin 100 mg and placebo, respectively).

For the subset of patients with an eGFR before treatment of 45 to < 60 mL/min/1.73 m², the incidence rates of adverse reactions related to volume depletion were similar: 2.3 events per 100 patient-years for canagliflozin 100 mg and 2.6 events per 100 patient-years of exposure for placebo. In the same study, for patients with an eGFR 30 to < 45 mL/min/1.73 m² the incidence rate was higher for canagliflozin 100 mg (4.9 events per 100 patient-years) than for placebo (2.6 events per 100 patient-years).

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Laboratory values, described below, are derived from the pooled analysis of 26-week, placebo-controlled clinical studies unless otherwise noted.

Increases in serum potassium: Mean percent changes from baseline in blood potassium were 0.5% and 1.0% for canagliflozin 100 mg and 300 mg, respectively, compared to 0.6% for placebo. Episodes of elevated serum potassium (> 5.4 mEq/L and 15% above baseline) were seen in 4.4% of patients treated with canagliflozin 100 mg, 7.0% of patients treated with canagliflozin 300 mg, and 4.8% of patients treated with placebo.

In a trial in patients with moderate renal impairment (eGFR 30 to < 50 mL/min/1.73 m²), increases in serum potassium to >5.4 mEq/L and 15% above baseline were seen in 16.1%, 12.4%, and 27.0% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Elevations to ≥6.5 mEq/L occurred in 1.1%, 2.2%, and 2.2% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

In a long-term renal outcomes study in patients with type 2 diabetes and diabetic nephropathy, no increase in adverse events of hyperkalemia, and no absolute (> 6.5 mEq/L) or relative (>upper limit of normal and > 15% increase from baseline) increases in serum potassium were observed with canagliflozin 100 mg relative to placebo.

Increases in serum creatinine and blood urea nitrogen (BUN): Mean percent changes from baseline in creatinine, with commensurate decreases in eGFR, were 2.8% and 4.0% for canagliflozin 100 mg and 300 mg, respectively, compared to 1.5% for placebo. Mean percent increases from baseline in BUN were 17.1% and 18.0% for canagliflozin 100 mg and 300 mg, respectively, compared to 2.7% for placebo. These changes were generally observed within six

weeks of treatment initiation. Subsequently, serum creatinine concentrations gradually trended toward baseline and BUN levels remained stable.

The proportion of patients with larger decreases in eGFR (> 30%) from baseline, occurring at any time during treatment, was 2.0% with canagliflozin 100 mg and 4.1% with canagliflozin 300 mg relative to 2.1% with placebo. At study end, decreases of > 30% from baseline were seen for 0.7% of subjects with canagliflozin 100 mg, 1.4% with canagliflozin 300 mg, and 0.5% with placebo (see [7 WARNINGS AND PRECAUTIONS](#)). After discontinuation of canagliflozin therapy, these changes in laboratory values improved or returned to baseline.

In an integrated analysis of data from two long-term cardiovascular outcome studies, patients treated with canagliflozin experienced an initial fall in mean eGFR that thereafter stabilized (see Figure 1) whereas patients treated with placebo experienced a progressive decline in eGFR.




Figure 1: Adjusted mean eGFR over time

In a long-term renal outcomes trial, patients treated with canagliflozin experienced an acute decrease in eGFR at Week 3, followed by an attenuated decline over time from week 3 to end of treatment. Placebo-treated patients demonstrated a progressive linear decline over time. After Week 52, the LS mean decrease in eGFR was smaller in the canagliflozin 100 mg group than in the placebo group (Figure 2).




Figure 2: LS Mean Change From Baseline in eGFR Over Time (On-Treatment Analysis Set)

Lipid changes: Compared to placebo, mean increases from baseline in low density lipoprotein cholesterol (LDL-C) were 0.11 mmol/L (4.5%) and 0.21 mmol/L (8.0%) with canagliflozin 100 mg and canagliflozin 300 mg, respectively. Increases in total cholesterol of 0.12 mmol/L (2.5%) and 0.21 mmol/L (4.3%) were seen, relative to placebo, for canagliflozin 100 mg and canagliflozin 300 mg, respectively. Increases in non-HDL-C relative to placebo were 0.05 mmol/L (1.5%) and 0.13 mmol/L (3.6%) with canagliflozin 100 mg and 300 mg, respectively. Increases in high-density lipoprotein cholesterol (HDL-C) were 0.06 mmol/L (5.4%), and 0.07 mmol/L (6.3%) relative to placebo for canagliflozin 100 mg and canagliflozin 300 mg, respectively. The LDL-C/HDL-C ratios did not change with either canagliflozin dose compared to placebo.

Increases in hemoglobin: Mean hemoglobin concentration increased from baseline 4.7 g/L (3.5%) with canagliflozin 100 mg and 5.1 g/L (3.8%) with canagliflozin 300 mg, compared to a decrease of -1.8 g/L (-1.1%) with placebo. After 26 weeks of treatment, 0.8%, 4.0%, and 2.7% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively, had a hemoglobin level above the upper limit of normal.

Increases in serum phosphate: Dose-related increases in serum phosphate levels were observed with canagliflozin. In the pool of four placebo-controlled trials, the mean percent change in serum phosphate levels were 3.6% and 5.1% with canagliflozin 100 mg and canagliflozin 300 mg, respectively, compared to 1.5% with placebo. Episodes of elevated serum phosphate (> 1.65 mmol/L and 25% above baseline) were seen in 0.6% and 1.6% of patients treated with canagliflozin 100 mg and 300 mg, respectively, compared to 1.3% of patients treated with placebo.

Decreases in serum urate: Moderate decreases in the mean percent change from baseline in serum urate were observed in the canagliflozin 100 mg and 300 mg groups (-10.1% and -10.6%, respectively) compared with placebo, where a slight increase from baseline (1.9%) was observed. Decreases in serum urate in the canagliflozin groups were maximal or near maximal

by Week 6 and maintained with dosing. A transient increase in urinary uric acid excretion was seen, which was not persistent.

Electrolytes: The following changes from baseline to end of treatment in serum electrolytes were observed during canagliflozin treatment in the CANVAS integrated database.

Table 11: Placebo-adjusted Mean Changes from Baseline in Electrolytes at Week 18 or 26^a in the CANVAS program

Analyte [normal range, unit]	Baseline, mean (SE)	Placebo-corrected change from baseline at Week 18 or 26 ^a , mean (95%)	p-value
Sodium [135 – 145 mmol/L]			
canagliflozin	139.3 (0.036)	0.40 (0.304;0.496)	<0.001
Potassium [3.5 – 5.0 mmol/L]			
canagliflozin	4.44 (0.006)	0.01 (-0.005;0.028)	0.171
Magnesium [0.75 – 0.95 mmol/L]			
canagliflozin	0.77 (0.001)	0.08 (0.074; 0.080)	<0.001
Bicarbonate [24 – 30 mmol/L]			
canagliflozin	23.33 (0.036)	-0.41 (-0.504;-0.307)	<0.001
Phosphate [0.80-1.50 mmol/L]			
canagliflozin	1.16 (0.002)	0.03 (0.028;0.040)	<0.001
Calcium [2.07-2.64 mmol/L]			
canagliflozin	2.41 (0.002)	0.02 (0.012, 0.020)	<0.001

^a CANVAS study blood chemistries obtained at week 18, CANVAS-R study blood chemistries obtained at week 26

SE = standard error

ANCOVA for Week 18 or 26 includes the baseline electrolyte as a linear covariate, and treatment and study as fixed effects.

The following shifts from normal range at baseline to below or above the normal range at worst value on treatment were reported in the treated set in the CANVAS integrated database:

- Increases in serum sodium above the upper limit of normal occurred more frequently in patients receiving canagliflozin than in those receiving placebo (2.63 per 100 subject years for canagliflozin and 1.80 per 100 subject years for placebo).
- Decreases in serum magnesium below the lower limit of normal occurred more frequently in patients receiving placebo (0.65 per 100 subject years for canagliflozin and 3.80 per 100 subject years for placebo), whilst increases in serum magnesium above the upper limit of normal occurred more frequently in patients receiving canagliflozin than in those receiving placebo (1.25 per 100 subject years for canagliflozin and 0.88 per 100 subject years for placebo).
- Decreases of serum bicarbonate below the lower limit of normal occurred more frequently in patients receiving canagliflozin than in those receiving placebo (2.91 per 100 subject years for canagliflozin, 2.39 per 100 subject years for placebo).
- Increases of serum phosphate above the upper limit of normal occurred more frequently in patients receiving canagliflozin than in those receiving placebo (1.36 per 100 subject years for canagliflozin and 1.00 per 100 subject years for placebo).

8.5 Post-Market Adverse Reactions

Canagliflozin

Gastrointestinal Disorders: pancreatitis acute

Metabolism and nutrition disorders: diabetic ketoacidosis

Immune system disorders: anaphylactic reaction

Skin and subcutaneous tissue disorders: angioedema

Renal and urinary disorders: acute kidney injury, including acute renal failure (with or without volume depletion).

Genitourinary: severe urinary tract infections; urosepsis and pyelonephritis

Musculoskeletal: bone fractures

Infections and Infestations: Fournier's gangrene (necrotizing fasciitis of the perineum)

Metformin hydrochloride

Blood and Lymphatic System Disorders: Hemolytic anemia, some with a fatal outcome

Gastrointestinal Disorders: Abdominal discomfort, abdominal distension, abdominal pain, abdominal pain upper, constipation, diarrhea, dry mouth, dyspepsia, flatulence, gastric disorder, gastric ulcer, gastrointestinal disorder, nausea, vomiting

Hepatobiliary Disorders: Liver function tests abnormalities or hepatitis resolving upon metformin discontinuation, autoimmune hepatitis, drug-induced liver injury, hepatitis

Investigations: Blood lactic acid increased

Metabolism and Nutrition Disorders: Lactic acidosis, decrease of Vitamin B₁₂ absorption with decrease of serum levels during long-term use of metformin, weight decreased, decreased appetite

Nervous System Disorders: Encephalopathy, peripheral neuropathy in patients with vitamin B₁₂ deficiency

Skin and Subcutaneous Tissue Disorders: Erythema, pruritus, rash, skin lesion, urticaria

9 DRUG INTERACTIONS

9.2 Drug Interactions Overview

Specific pharmacokinetic drug interactions studies with INVOKAMET have not been performed, although such studies have been conducted with the individual canagliflozin and metformin components.

Co-administration of canagliflozin (300 mg once daily) and metformin (2000 mg once daily) had no clinically relevant effect on the pharmacokinetics of either canagliflozin or metformin.

Canagliflozin

Canagliflozin did not induce CYP450 enzyme expression (3A4, 2C9, 2C19, 2B6, and 1A2) in cultured human hepatocytes. Canagliflozin did not inhibit the CYP450 isoenzymes (1A2, 2A6, 2C19, 2D6, or 2E1) and weakly inhibited CYP2B6, CYP2C8, CYP2C9, and CYP3A4 based on *in vitro* studies with human hepatic microsomes. Canagliflozin is a weak inhibitor of P-gp.

Canagliflozin is also a substrate of drug transporters P-glycoprotein (P-gp), Breast Cancer Resistance Protein (BCRP) and Multi-Drug Resistance-Associated Protein 2 (MRP2).

Inhibition of BCRP by canagliflozin cannot be excluded at an intestinal level and increased exposure may therefore occur for drugs transported by BCRP, e.g., certain statins like rosuvastatin and some anti-cancer agents.

9.3 Drug-Behavioural Interactions

Patients should be alerted to the elevated risk of adverse reactions related to reduced intravascular volume, such as postural dizziness, and to the risk of hypoglycemia when INVOKAMET is used as add-on therapy with insulin or an insulin secretagogue (see [7 WARNINGS AND PRECAUTIONS](#), [8 ADVERSE REACTIONS](#) and [4.1 Dosing Considerations](#)).

Patients should be cautioned against excessive alcohol intake, either acute or chronic, when taking INVOKAMET, since alcohol intake potentiates the effect of metformin on lactate metabolism (see [2 CONTRAINDICATIONS](#)). The risk of lactic acidosis is increased in acute alcohol intoxication, particularly in case of fasting or malnutrition or hepatic insufficiency. INVOKAMET is contraindicated in patients with clinical or laboratory evidence of hepatic disease (see [2 CONTRAINDICATIONS](#)). It is recommended that consumption of alcohol and alcohol-containing medicinal product be avoided.

9.4 Drug-Drug Interactions

The drugs listed are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction.

Effects of other drugs on canagliflozin

In clinical studies, the effects of other drugs on canagliflozin were assessed. Cyclosporin (P-gp inhibitor), hydrochlorothiazide, oral contraceptives (ethinyl estradiol and levonorgestrel), metformin, and probenecid (UGT, MRP2, OATP, OAT1 and OAT3 inhibitor) had no clinically relevant effect on the pharmacokinetics of canagliflozin.

Table 12: Effect of Co-administered Drugs on Systemic Exposure of Canagliflozin

Co-administered Drug	Dose of Co-administered Drug ¹	Dose of Canagliflozin ¹	Geometric Mean Ratio (Ratio With/Without Co-administered Drug) No Effect = 1.0		Clinical Comment
			AUC ² (90% CI)	C _{max} (90% CI)	
Cyclosporin	400 mg	300 mg once daily for 8 days	1.23 (1.19; 1.27)	1.01 (0.91; 1.11)	No dosage adjustment for INVOKAMET required
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg once daily for 6 days	0.91 (0.88; 0.94)	0.92 (0.84; 0.99)	No dosage adjustment for INVOKAMET required
Hydrochlorothiazide	25 mg once daily for 35 days	300 mg once daily for 7 days	1.12 (1.08; 1.17)	1.15 (1.06; 1.25)	No dosage adjustment for INVOKAMET required
Probenecid	500 mg twice daily for 3 days	300 mg once daily for 17 days	1.21 (1.16; 1.25)	1.13 (1.00; 1.28)	No dosage adjustment for INVOKAMET required
Inducers of UGT enzymes / drug transporters					
Rifampin	600 mg once daily for 8 days	300 mg	0.49 (0.44; 0.54)	0.72 (0.61; 0.84)	Consider increasing the INVOKAMET dose to 300 mg once daily if patients are currently tolerating INVOKAMET 100 mg once daily (refer to 4 DOSAGE AND ADMINISTRATION).
Phenytoin, phenobarbital, barbiturates, carbamazepine, ritonavir, efavirenz, or St. John's Wort	N/A ³				Consider increasing the INVOKAMET dose to 300 mg once daily if patients are currently tolerating INVOKAMET 100 mg once daily (refer to 4 DOSAGE AND ADMINISTRATION).

¹ Single dose unless otherwise noted

² AUC_{inf} for drugs given as a single dose and AUC_{24h} for drugs given as multiple doses.

³ N/A = Not applicable

Effects of canagliflozin on other drugs

Canagliflozin at steady-state had no clinically relevant effect on the pharmacokinetics of metformin, oral contraceptives (ethinyl estradiol and levonorgestrel-CYP3A4 substrates), glyburide (CYP2C9 substrate), simvastatin (CYP3A4 substrate), acetaminophen, hydrochlorothiazide, or warfarin (CYP2C9 substrate), in healthy subjects.

Pharmacokinetic Interactions

Lithium: The concomitant use of INVOKAMET or other SGLT2 inhibitors with lithium may decrease serum lithium levels through increased renal lithium elimination. Therefore, serum lithium concentration should be monitored more frequently with INVOKAMET initiation, following dosage changes or following discontinuation. The patient should be referred to their lithium prescriber for serum lithium concentration monitoring and clinical supervision as required during treatment.

Table 13: Effect of Canagliflozin on Systemic Exposure of Co-Administered Drugs

Co-Administered Drug	Dose of Co-Administered Drug ¹	Dose of Canagliflozin ¹	Geometric Mean Ratio (Ratio With/Without Co-Administered Drugs) No Effect = 1.0			Clinical Comment
				AUC ² (90% CI)	C _{max} (90% CI)	
Digoxin	0.5 mg once daily first day followed by 0.25 mg once daily for 6 days	300 mg once daily for 7 days	Digoxin	1.20 (1.12; 1.28)	1.36 (1.21; 1.53)	Patients taking INVOKAMET with concomitant digoxin should be monitored appropriately
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg once daily for 6 days	Ethinyl estradiol	1.07 (0.99; 1.15)	1.22 (1.10; 1.35)	No dosage adjustment required for ethinyl estradiol and levonorgestrel
			Levonorgestrel	1.06 (1.00; 1.13)	1.22 (1.11; 1.35)	
Glyburide	1.25 mg	200 mg once daily for 6 days	Glyburide	1.02 (0.98; 1.07)	0.93 (0.85; 1.01)	No dosage adjustment required for glyburide
			3-cis-hydroxy-glyburide	1.01 (0.96; 1.07)	0.99 (0.91; 1.08)	
			4-trans-hydroxy-glyburide	1.03 (0.97; 1.09)	0.96 (0.88; 1.04)	
Hydrochlorothiazide	25 mg once daily for 35 days	300 mg once daily for 7 days	Hydrochlorothiazide	0.99 (0.95; 1.04)	0.94 (0.87; 1.01)	No dosage adjustment required for hydrochlorothiazide
Acetaminophen	1000 mg	300 mg twice daily for 25 days	Acetaminophen	1.06 ³ (0.98; 1.14)	1.00 (0.92; 1.09)	No dosage adjustment required for acetaminophen
Simvastatin	40 mg	300 mg once daily for 7 days	Simvastatin	1.12 (0.94; 1.33)	1.09 (0.91; 1.31)	No dosage adjustment required for simvastatin
			Simvastatin acid	1.18 (1.03; 1.35)	1.26 (1.10; 1.45)	
Warfarin	30 mg	300 mg once daily for 12 days	(R)-warfarin	1.01 (0.96; 1.06)	1.03 (0.94; 1.13)	No dosage adjustment

Co-Administered Drug	Dose of Co-Administered Drug ¹	Dose of Canagliflozin ¹	Geometric Mean Ratio (Ratio With/Without Co-Administered Drugs) No Effect = 1.0			Clinical Comment
				AUC ² (90% CI)	C _{max} (90% CI)	
			(S)-warfarin	1.06 (1.00; 1.12)	1.01 (0.90; 1.13)	required for warfarin

¹ Single dose unless otherwise noted

² AUC_{inf} for drugs given as a single dose and AUC_{24h} for drugs given as multiple doses.

³ AUC_{0-12h}

Metformin hydrochloride

Glyburide: In a single-dose interaction study in type 2 diabetes patients, coadministration of metformin and glyburide did not result in any changes in either metformin pharmacokinetics or pharmacodynamics. Decreases in glyburide AUC and C_{max} were observed, but were highly variable. The single-dose nature of this study and the lack of correlation between glyburide blood levels and pharmacodynamic effects make the clinical significance of this interaction uncertain.

Furosemide: A single-dose, metformin-furosemide drug interaction study in healthy subjects demonstrated that pharmacokinetic parameters of both compounds were affected by coadministration. Furosemide increased the metformin plasma and blood C_{max} by 22% and blood AUC by 15%, without any significant change in metformin renal clearance. When administered with metformin, the C_{max} and AUC of furosemide were 31% and 12% smaller, respectively, than when administered alone, and the terminal half-life was decreased by 32%, without any significant change in furosemide renal clearance. No information is available about the interaction of metformin and furosemide when coadministered chronically.

Nifedipine: A single-dose, metformin-nifedipine drug interaction study in normal healthy volunteers demonstrated that coadministration of nifedipine increased plasma metformin C_{max} and AUC by 20% and 9%, respectively, and increased the amount excreted in the urine. T_{max} and half-life were unaffected. Nifedipine appears to enhance the absorption of metformin. Metformin had minimal effects on nifedipine.

Cationic drugs: Cationic drugs (e.g., amiloride, digoxin, morphine, procainamide, quinidine, quinine, ranitidine, triamterene, trimethoprim, or vancomycin) that are eliminated by renal tubular secretion, theoretically have the potential for interaction with metformin by competing for common renal tubular transport systems. Such an interaction has been observed between metformin and oral cimetidine in normal healthy volunteers in both single- and multiple-dose, metformin-cimetidine drug interaction studies, with a 60% increase in peak metformin plasma and whole blood concentrations and a 40% increase in plasma and whole blood metformin AUC.

There was no change in elimination half-life in the single-dose study. Metformin had no effect on cimetidine pharmacokinetics. Therefore, careful patient monitoring and dose adjustment of

metformin or the interfering drug is recommended in patients who are taking cationic medications that are excreted via renal tubular secretion.

Anticoagulant: Elimination rate of the anticoagulant phenprocoumon has been reported to be increased by 20% when used concurrently with metformin. Therefore, patients receiving phenprocoumon or other antivitamin K anticoagulants should be monitored carefully when both types of drugs are used simultaneously. In such cases, an important increase of prothrombin time may occur upon cessation of INVOKAMET therapy, with an increased risk of hemorrhage.

Other: Other drugs tend to produce hyperglycemia and may lead to loss of blood sugar control. These include the thiazides and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, estrogen plus progestogen, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blocking drugs, isoniazid and beta-2-agonists. ACE inhibitors may decrease the blood glucose levels. When such drugs are administered to a patient receiving metformin the patient should be closely observed to maintain adequate glycemic control. Diuretics, especially loop diuretics, may increase the risk of lactic acidosis due to their potential to decrease renal function.

INVOKAMET is not recommended for use in patients receiving loop diuretics. Canagliflozin may add to the effect of diuretics and may increase the risk of hypovolemia and hypotension (see [7 WARNINGS AND PRECAUTIONS](#) and [4 DOSAGE AND ADMINISTRATION](#)).

9.5 Drug-Food Interactions

Interactions with food have not been established.

9.6 Drug-Herb Interactions

St John's Wort (*Hypericum perforatum*) is a CYP3A4 inducer and co-administration with INVOKAMET may result in loss of efficacy or reduced clinical response. Dosage adjustment may be required (see [4.2 Recommended Dose and Dosage Adjustment](#)).

9.7 Drug-Laboratory Test Interactions

Due to its mechanism of action, patients taking INVOKAMET will test positive for glucose in their urine.

Increases in urinary glucose excretion with INVOKAMET can falsely lower 1,5-anhydroglucitol (1,5 AG) levels and make measurements of 1,5 AG unreliable in assessing glycemic control. Therefore, 1,5-AG assays should not be used for assessment of glycemic control in patients on INVOKAMET. For further detail, it may be advisable to contact the specific manufacturer of the 1,5-AG assay.

Intravascular contrast studies with iodinated materials can lead to acute alteration of renal function and have been associated with lactic acidosis in patients receiving metformin (see [2 CONTRAINDICATIONS](#) and [7 WARNINGS AND PRECAUTIONS, Renal](#)).

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

INVOKAMET (canagliflozin/metformin hydrochloride) combines two oral antihyperglycemic agents with complementary mechanisms of action to improve glycemic control in patients with type 2 diabetes: canagliflozin, a sodium-glucose co-transporter 2(SGLT2) inhibitor, and metformin hydrochloride, a member of the biguanide class.

Canagliflozin

Sodium-glucose co-transporter 2 (SGLT2), expressed in the proximal renal tubules, is responsible for the majority of the reabsorption of filtered glucose from the tubular lumen. Patients with diabetes have been shown to have elevated renal glucose reabsorption which may contribute to persistent elevated glucose concentrations. Canagliflozin is an orally-active inhibitor of SGLT2. By inhibiting SGLT2, canagliflozin reduces reabsorption of filtered glucose and lowers the renal threshold for glucose (RT_G), and thereby increases urinary glucose excretion, which decreases elevated plasma glucose concentrations by an insulin-independent mechanism in patients with type 2 diabetes. The increased urinary glucose excretion with SGLT2 inhibition also translates to an osmotic diuresis, with the diuretic effect leading to a reduction in systolic blood pressure; the increase in urinary glucose excretion results in a loss of calories and therefore a reduction in body weight, as demonstrated in studies of patients with type 2 diabetes.

Canagliflozin's action to increase UGE directly lowering plasma glucose is independent of insulin. Improvement in homeostasis model assessment for beta-cell function (HOMA beta-cell) and improved beta-cell insulin secretion response to a mixed-meal challenge has been observed in clinical studies with canagliflozin.

In Phase 3 studies, pre-meal administration of canagliflozin 300 mg provided a greater reduction in post-meal glucose excursion than observed with the 100 mg dose. This effect at the 300 mg dose of canagliflozin may, in part, be due to local inhibition of intestinal SGLT1 (an important intestinal glucose co-transporter) related to transient high concentrations of canagliflozin in the intestinal lumen prior to drug absorption (canagliflozin is a low potency inhibitor of SGLT1). Studies have shown no glucose malabsorption with canagliflozin.

Canagliflozin increases the delivery of sodium to the distal tubule by blocking SGLT2-dependent glucose and sodium reabsorption.

Metformin hydrochloride

Metformin is a biguanide derivative producing an antihyperglycemic effect which can only be observed in man or in the diabetic animal and only when there is insulin secretion. Metformin, at therapeutic doses, does not cause hypoglycemia when used alone in man or in the non-diabetic animal, except when using a near lethal dose. Metformin has no effects on the pancreatic beta cells. The mode of action of metformin is not fully understood. It has been postulated that metformin might potentiate the effect of insulin or that it might enhance the effect of insulin on the peripheral receptor site. This increased sensitivity seems to follow an increase in the number of insulin receptors on cell surface membranes.

10.2 Pharmacodynamics

Following single and multiple oral doses of canagliflozin to patients with type 2 diabetes, dose-dependent decreases in RT_G and increases in urinary glucose excretion were observed. From a starting value of RT_G of approximately 13 mmol/L, maximal suppression of 24-hour mean RT_G was seen with the 300 mg daily dose to approximately 4 to 5 mmol/L in patients with type 2 diabetes in Phase 1 studies (see model in Figure 3), suggesting a low risk for treatment-induced hypoglycemia. The reductions in RT_G led to increased UGE in subjects with type 2 diabetes treated with either 100 mg or 300 mg of canagliflozin ranging from 77 to 119 g/day across the Phase 1 studies; the UGE observed translates to a loss of 308 to 476 kcal/day. The reductions in RT_G and increases in UGE were sustained over a 26-week dosing period in patients with type 2 diabetes. Moderate increases (generally < 400-500 mL) in daily urine volume were seen that attenuated over several days of dosing. Urinary uric acid excretion was transiently increased by canagliflozin (increased by 19% compared to baseline on day 1 and then attenuating to 6% on day 2 and 1% on day 13). This was accompanied by a sustained reduction in serum uric acid concentration of approximately 20%.




Figure 3: Predicted (PK/PD Modelled) 24-Hour Profile for RT_G in Subjects with Type 2 Diabetes Treated with Canagliflozin 100 mg and 300 mg

In a single-dose study in patients with type 2 diabetes, treatment with 300 mg before a mixed meal delayed intestinal glucose absorption and reduced postprandial glucose through both renal and non-renal mechanisms.

Cardiac electrophysiology

In a randomized, double-blind, placebo-controlled, active-comparator, 4-way crossover study, 60 healthy subjects were administered a single oral dose of canagliflozin 300 mg, canagliflozin 1200 mg (4 times the maximum recommended dose), moxifloxacin, and placebo. No meaningful changes in QT_c interval were observed with either the recommended dose of 300 mg or the 1200 mg dose. At the 1200 mg dose, peak canagliflozin plasma concentrations were approximately 1.4 times the steady-state peak concentrations following a 300 mg once-daily

dose.

10.3 Pharmacokinetics

Pharmacokinetics of canagliflozin were comparable between healthy volunteers and type 2 diabetic patients based on clinical trials and population pharmacokinetic data. After single-dose oral administration of 100 mg and 300 mg in healthy subjects, canagliflozin was rapidly absorbed, with peak plasma concentrations (median T_{max}) occurring 1 to 2 hours post-dose. Plasma C_{max} and AUC of canagliflozin increased in a dose-proportional manner from 50 mg to 300 mg. The apparent terminal half-life ($t_{1/2}$) (expressed as mean \pm standard deviation) was 10.6 ± 2.13 hours to 13.1 ± 3.28 hours for the 100 mg and 300 mg doses, respectively. Steady-state was reached after 4 to 5 days of once-daily dosing with canagliflozin 100 mg to 300 mg. Canagliflozin does not exhibit time-dependent pharmacokinetics, and accumulated in plasma up to 36% following multiple doses of 100 mg and 300 mg.

Table 14: Summary of Canagliflozin’s Pharmacokinetic Parameters in Healthy Subjects and T2DM Patients at Steady State

	N	C_{max} (SD) (ng/mL)	$t_{1/2}$ (h)	AUC _{24h} (SD) (ng.h/mL)	Cl/F	Vd/F
Healthy Volunteers^a						
100 mg multiple oral doses qd	9	1,118 (143)	13.3 (4.8)	6,056 (959)	16.4 (2.16)	304 (79.7)
300 mg multiple oral doses qd	9	3,379 (728)	13.5 (3.2)	19,252 (5,348)	16.4 (3.60)	319 (104)
T2DM Patients^b						
100 mg multiple oral doses qd	8	1,227 (481)	13.7 (2.1)	8,225 (1,947)	13.0 (4.43)	250 (50.7)
300 mg multiple oral doses qd	10	4,678 (1,685)	14.9 (4.8)	30,995 (11,146)	11.3 (5.21)	226 (89.4)

^a From Study DIA1030

^b From Study DIA1023

The results of bioequivalence studies in healthy subjects demonstrated that INVOKAMET 50 mg/500 mg, 50 mg/1000 mg, 150 mg/500 mg, and 150 mg/1000 mg combination tablets are bioequivalent to co-administration of corresponding doses of canagliflozin and metformin hydrochloride as individual tablets.

Administration of INVOKAMET 150 mg/1000 mg fixed-dose combination with food resulted in no change in overall exposure of canagliflozin. There was no change in metformin AUC; however, mean peak plasma concentration of metformin was decreased by 16% when administered with food. A delayed time to peak plasma concentration was observed for both components (2 hours for canagliflozin and 1 hour for metformin) under fed conditions. These changes are not likely to be clinically significant. As metformin is recommended to be administered with a meal to reduce the incidence of gastrointestinal side effects, it is recommended that INVOKAMET be taken with a meal to reduce gastrointestinal intolerability associated with metformin.

The 24-h mean RTG at steady state was similar following once daily and twice daily dosing regimens at the same total daily dose of 100 mg or 300 mg. The mean systemic exposure (AUC) at steady state was similar following once daily and twice daily dosing regimens at the same total daily dose of 100 mg or 300 mg.

Absorption

Canagliflozin

The mean absolute oral bioavailability of canagliflozin is approximately 65%. Co-administration of a high-fat meal with canagliflozin had no effect on the pharmacokinetics of canagliflozin; therefore, canagliflozin may be taken with or without food (see [4.4 Administration](#)).

Metformin hydrochloride

Metformin absorption is relatively slow and may extend over 6 hours.

Distribution

Canagliflozin

The mean steady-state volume of distribution of canagliflozin following a single intravenous infusion in healthy subjects was 119 L, suggesting extensive tissue distribution. Canagliflozin is extensively bound to proteins in plasma (99%), mainly to albumin. Protein binding is independent of canagliflozin plasma concentrations. Plasma protein binding is not meaningfully altered in patients with renal or hepatic impairment.

Metformin hydrochloride

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean V_d ranged between 63 – 276 liters.

Metabolism

Canagliflozin

O-glucuronidation is the major metabolic elimination pathway for canagliflozin, which is mainly glucuronidated by UDP glucuronosyl transferase 1A9 (UGT1A9) and 2B4 (UGT2B4) to two inactive O-glucuronide metabolites. CYP3A4-mediated (oxidative) metabolism of canagliflozin is minimal (approximately 7%) in humans.

Metformin hydrochloride

Metformin is not metabolized. Its main sites of concentration are the intestinal mucosa and the salivary glands. The plasma concentration at steady-state ranges about 1 to 2 mcg/mL. Certain drugs may potentiate the effects of metformin (see [7 WARNINGS AND PRECAUTIONS](#) and [9.4 Drug-Drug Interactions](#)).

Elimination

Canagliflozin

Following administration of a single oral [14 C] canagliflozin dose to healthy subjects, 41.5%, 7.0%, and 3.2% of the administered radioactive dose was recovered in faeces as canagliflozin, a hydroxylated metabolite, and an O-glucuronide metabolite, respectively. Enterohepatic circulation of canagliflozin was negligible.

Approximately 33% of the administered radioactive dose was excreted in urine, mainly as O-glucuronide metabolites (30.5%). Less than 1% of the dose was excreted as unchanged

canagliflozin in urine. Renal clearance for the 100 mg and 300 mg doses ranged from 1.30 to 1.55 mL/min.

Canagliflozin is a low-clearance drug, with a mean systemic clearance of approximately 192 mL/min in healthy subjects following intravenous administration.

Metformin hydrochloride

Metformin is excreted in urine at high renal clearance rate of about 450 mL/min. The initial elimination of metformin is rapid with a half-life varying between 1.7 and 3 hours. The terminal elimination phase accounting for about 4 to 5 % of the absorbed dose is slow with a half-life between 9 and 17 hours.

Special Populations and Conditions

- **Pediatrics:** Based on the data submitted and reviewed by Health Canada, the safety and efficacy of canagliflozin in pediatric patients <18 years of age have not been established; therefore, Health Canada has not authorized an indication for pediatric use (see [7.1.3 Pediatrics](#)).

An open-label, sequential, multiple-dose, multicentre pediatric Phase 1 study examined the pharmacokinetics and pharmacodynamics of canagliflozin in children and adolescents ≥11 to <18 years of age (mean age 14.6 years) with type 2 diabetes mellitus who were on a stable dose of metformin. The mean body weight was 107.15 kg (range: 48.5 to 168.6 kg).

The patients were treated with canagliflozin once-daily 100 mg or 300 mg for 14 days.

Table 15: Mean (SD) Plasma Canagliflozin Pharmacokinetic Parameters on Day 14

Parameters	Canagliflozin 100 mg QD (N=8) Mean (Std. Dev.)	Canagliflozin 300 mg QD (N=9) Mean (Std. Dev.)
C _{max} (ng/mL)	951 (429)	3,260 (1,330)
AUC (h*ng/mL)	6,190 (1,770)	28,392 (12,412)
t _{1/2} (h)	11.3 (2.5)	15.2 (6.9)
CL _{ss} /F (L/h)	17.5 (5.78)	12.3 (6.90)

- **Geriatrics:** Studies characterizing the pharmacokinetics of canagliflozin and metformin after administration of INVOKAMET in geriatric patients have not been performed.

Canagliflozin

Age had no clinically meaningful effect on the pharmacokinetics of canagliflozin based on a population pharmacokinetic analysis. However, patients 65 years and older had a higher incidence of adverse reactions related to reduced intravascular volume with INVOKAMET (see [7 WARNINGS AND PRECAUTIONS](#), [8 ADVERSE REACTIONS](#) and [4 DOSAGE AND ADMINISTRATION](#)).

Metformin hydrochloride

Controlled clinical studies of metformin did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients.

- **Sex:** Dose normalized exposures of canagliflozin in females were 22% higher than males, based on population pharmacokinetic analysis. These increases in exposures are not clinically meaningful and hence no dosage adjustment of canagliflozin is necessary based on sex.
- **Genetic Polymorphism:** Both UGT1A9 and UGT2B4 are subject to genetic polymorphism. In a pooled analysis of clinical data, increases in canagliflozin AUC of 26% were observed in UGT1A9*1/*3 carriers and 18% in UGT2B4*2/*2 carriers. These increases in canagliflozin exposure are not expected to be clinically relevant and no dosage adjustment is necessary based on UGT1A9 and UGT2B4 genetic polymorphisms. The effect of being homozygote (UGT1A9*3/*3, frequency < 0.1%) is probably more marked, but has not been investigated.
- **Ethnic Origin:** Dose normalized exposures of canagliflozin were comparable in white and non-white subjects, Blacks, Asians, and other ethnic origins. A population PK analysis of canagliflozin in 942 white subjects and 674 non-white subjects showed no significant impact of ethnic origin on canagliflozin PK and hence no dosage adjustment of canagliflozin is necessary based on ethnic origin.

No studies of metformin pharmacokinetic parameters according to ethnic origin have been performed.

- **Hepatic Insufficiency:**

Canagliflozin

INVOKAMET is contraindicated in patients with clinical or laboratory evidence of hepatic disease (see [2 CONTRAINDICATIONS](#)). Relative to subjects with normal hepatic function, the geometric mean ratios for C_{max} and AUC_{∞} of canagliflozin were 107% and 110%, respectively, in subjects with Child-Pugh class A (mild hepatic impairment) and 96% and 111%, respectively, in subjects with Child-Pugh class B (moderate hepatic impairment) following administration of a single 300 mg dose of canagliflozin.

These differences are not considered to be clinically meaningful. There is no clinical experience in patients with Child-Pugh class C (severe) hepatic impairment and therefore, canagliflozin is not recommended for use in this patient population.

Metformin hydrochloride

No pharmacokinetic studies of metformin have been conducted in patients with hepatic insufficiency.

- **Renal Insufficiency:**

Canagliflozin

A single-dose, open-label study evaluated the pharmacokinetics of canagliflozin 200 mg in subjects with varying degrees of renal impairment, classified using the Modification of Diet in Renal Disease (MDRD)-eGFR formula, compared to healthy subjects. The study included 3 subjects with normal renal function (eGFR ≥ 90 mL/min/1.73 m²), 10 subjects with mild renal impairment (eGFR 60 to <90 mL/min/1.73 m²), 9 subjects with moderate renal impairment (eGFR 30 to <60 mL/min/1.73 m²), and 10 subjects with severe renal impairment (eGFR 15 to <30 mL/min/1.73 m²) as well as 8 subjects with end stage renal disease (ESRD) on hemodialysis.

The C_{max} of canagliflozin was moderately increased by 13%, 29%, and 29% in subjects with mild, moderate, and severe renal failure, respectively, but not in subjects on hemodialysis.

Compared to healthy subjects, plasma AUC of canagliflozin was increased by approximately 17%, 63%, and 50% in subjects with mild, moderate, and severe renal impairment, respectively, but was similar for ESRD subjects and healthy subjects. Increases in canagliflozin AUC of this magnitude are not considered clinically relevant, however, the pharmacodynamic response to canagliflozin declines with increasing severity of renal impairment (see [2 CONTRAINDICATIONS](#) and [7 WARNINGS AND PRECAUTIONS](#)). Canagliflozin was negligibly removed by hemodialysis.

- **Obesity:** For subjects with body weight <78.2 kg, the dose normalized exposures of canagliflozin increased by 33%, based on population pharmacokinetic analysis. These increases in exposures are not clinically meaningful and hence no dosage adjustment of canagliflozin is necessary based on body weight.

11 STORAGE, STABILITY AND DISPOSAL

INVOKAMET tablets should be stored at 15-30°C. Patients should be instructed to talk to their pharmacist about any medications that have expired, or that they no longer use. Medications should not be disposed in wastewater or household waste.



12 SPECIAL HANDLING INSTRUCTIONS

Keep INVOKAMET out of the sight and reach of children.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance: Canagliflozin plus metformin hydrochloride

Proper name:	Canagliflozin	Metformin hydrochloride
Chemical name:	(1S)-1,5-anhydro-1-[3-[[5-(4-fluorophenyl)-2-thienyl]methyl]-4-methylphenyl]-D-glucitol hemihydrate	1,1-Dimethylbiguanide hydrochloride
Molecular formula:	$C_{24}H_{25}FO_5S \cdot 1/2 H_2O$	$C_4H_{11}N_5 \cdot HCl$
Molecular mass:	<ul style="list-style-type: none">• Hemihydrate: 453.53• Anhydrous: 444.52	165.62
Structural formula:		
Physicochemical properties:	Canagliflozin is practically insoluble in aqueous media from pH 1.1 to 12.9. There is no detectable pK_a value for this substance.	Metformin hydrochloride is freely soluble in various aqueous media, irrespective of pH, and has low permeability.

14 CLINICAL TRIALS

14.1 Clinical Trials by Indication

Type 2 Diabetes Mellitus

Table 16: Summary of Patient Demographics for Clinical Trials in Type 2 Diabetes Mellitus

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Mean age (Range)	Sex (%F/M)
Add-on Therapy with Metformin (\geq 1500 mg/day)					
DIA3006	Randomized, double-blind, active-controlled, parallel-group, multicentre	Canagliflozin 100 or 300 mg/day or Sitagliptin 100 mg/day or Placebo 26-week	Total: 1284 Canagliflozin 100 mg: 368 Canagliflozin 300 mg: 367 Sitagliptin 100 mg: 366 Placebo: 183	55.4 (21-79)	52.9/47.1
DIA3009	Randomized, double-blind, active-controlled, parallel-group, multicentre	Canagliflozin 100 or 300 mg/day or Glimepiride 1- 8 mg (titration protocol) 52-week	Total: 1450 Canagliflozin 100 mg: 483 Canagliflozin 300 mg: 485 Glimepiride: 482	56.2 (22-80)	47.9/52.1
Add-on Therapy with Metformin (\geq 1500 mg/day) and a Sulfonylurea (stable dose)					
DIA3002	Randomized, double-blind, placebo-controlled, parallel-group, multicentre	Canagliflozin 100 or 300 mg/day or Placebo 26-week	Total: 469 Canagliflozin 100 mg: 157 Canagliflozin 300 mg: 156 Placebo: 156	56.8 (27-79)	49.0/51.0
DIA3015	Randomized, double-blind, active-controlled, parallel-group, multicentre	Canagliflozin 300 mg/day or Sitagliptin 100 mg/day or Placebo 52-week	Total: 755 Canagliflozin 300 mg: 377 Sitagliptin 100 mg: 378	56.7 (21-91)	44.1/55.9
Add-on Therapy with Metformin (\geq 1500 mg/day) and Pioglitazone (30 or 45 mg/day)					
DIA3012	Randomized, double-blind, placebo-controlled, parallel-group, multicentre	Canagliflozin 100 or 300 mg/day or Placebo 26-week	Total: 342 Canagliflozin 100 mg: 113 Canagliflozin 300 mg: 114 Placebo: 115	57.4 (27-78)	36.8/63.2

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Mean age (Range)	Sex (%F/M)
Cardiovascular					
DIA3008	Randomized, double-blind, placebo-controlled, parallel-group, Multicentre	Canagliflozin 100 or 300 mg/day or Placebo mean 223 weeks exposure to study drug	Total: 4330 Canagliflozin 100 mg: 1445 Canagliflozin 300 mg: 1443 Placebo: 1442	63 (32-87)	33.9/66.1
DIA4003	Randomized, double-blind, placebo-controlled, parallel-group, Multicentre	Canagliflozin 100 up-titrated to 300 mg/day at week 13 or later at investigators' discretion mean 94 weeks exposure to study drug	Total: 5813 Canagliflozin 100 mg uptitrated: 2907 Placebo: 2906	64 (30-89)	37.2/62.8
Renal					
DNE3001	Randomized, double-blind, placebo-controlled, parallel-group, Multicentre	Canagliflozin 100 mg or Placebo mean 115 weeks exposure to study drug	Total: 4401 Canagliflozin 100 mg: 2202 Placebo: 2199	63 (30-89)	33.9/66.1
Add-on to Insulin and Metformin					
DIA 3008 Insulin Substudy (subset)	Randomized, double-blind, placebo-controlled, parallel-group, Multicentre	Canagliflozin 100 or 300 mg/day or Placebo 18 weeks	Total: 731 Canagliflozin 100 mg: 241 Canagliflozin 300 mg: 246 Placebo: 244	57.0 (21-91)	46.1/53.9

¹ AHA = antihyperglycemic agent

There have been no clinical efficacy studies conducted with INVOKAMET (canagliflozin/metformin hydrochloride), however, bioequivalence of INVOKAMET to canagliflozin and metformin co-administered as individual tablets was demonstrated in healthy subjects (see [10.3 Pharmacokinetics](#)). The co-administration of canagliflozin and metformin has been studied in patients with type 2 diabetes inadequately controlled on diet and exercise and as add-on therapy with other antihyperglycemic agents.

In the metformin add-on studies, a total of 5,031 patients with type 2 diabetes were randomized in six double-blind, controlled clinical efficacy and safety studies conducted to evaluate the effects of canagliflozin on glycemic control. The racial distribution was 71% White, 15% Asian, 5% Black, and 9% other groups. Approximately 20% of patients were Hispanic. Approximately 54% of patients were male. Patients had an overall mean age of 59.6 years (range 21 to 91

years), with 1036 patients 65 years of age and older and 121 patients 75 years of age and older.

In addition, an 18-week double-blind, placebo-controlled Phase 2 study with twice daily dosing (canagliflozin 50 mg or 150 mg in combination with metformin) was conducted in 279 patients in which 186 patients were treated with canagliflozin in combination with metformin.

Study Results

In patients with type 2 diabetes, treatment with canagliflozin produced statistically significant improvements in HbA1c, fasting plasma glucose (FPG), 2-hour postprandial glucose (PPG), and body weight, compared to placebo. Canagliflozin was effective in reducing HbA1c in a broad range of patients regardless of disease duration and concomitant use of antihyperglycemic agents. The durability of these reductions in HbA1c was demonstrated in two Phase 3 studies, with minimal attenuation of the glycemic response to canagliflozin over 52 weeks, in contrast to the deterioration of the glycemic response observed with comparators.

Statistically significant improvements in glycemic control relative to placebo were observed with canagliflozin when given as add on therapy with metformin, add-on therapy with metformin and a sulfonylurea, add-on therapy with metformin and pioglitazone, or as add-on therapy with insulin and metformin).

In addition, significant improvements in HbA1c were observed with canagliflozin in older patients. Reductions in HbA1c were observed across subgroups including age, gender, ethnic origin, baseline body mass index (BMI), and baseline beta-cell function. Greater reductions in HbA1c relative to placebo were observed in patients with higher baseline HbA1c or eGFR values.

Add-on Therapy with Metformin (Study DIA3006)

A total of 1284 patients with inadequate glycemic control (HbA1c of $\geq 7\%$ to $\leq 10.5\%$) on metformin monotherapy (2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) participated in a randomized, double-blind, placebo- and active-controlled, parallel-group, 4-arm, multicentre clinical study to evaluate the efficacy of canagliflozin as add-on therapy with metformin over 26 weeks. The mean age was 55 years, 47% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m². Patients already on metformin (N=1009) at screening with inadequate glycemic control completed a 2-week, single-blind, placebo run-in period. Other patients on metformin and another oral agent or a lower than required dose of metformin (N=275) were switched to a regimen of metformin monotherapy. After at least 8 weeks on a stable dose of metformin monotherapy, patients entered a 2-week, single-blind, placebo run-in period. Patients were randomized to the addition of canagliflozin 100 mg, canagliflozin 300 mg, sitagliptin 100 mg, or placebo, administered once daily.

As shown in Table 17, statistically significant ($p < 0.001$) reductions in HbA1c, FPG, PPG, and body weight relative to placebo were observed. In addition, a greater percentage of patients achieved an HbA1c $< 7.0\%$ compared to placebo. Statistically significant ($p < 0.001$) reductions in systolic blood pressure were observed with canagliflozin 100 mg and 300 mg relative to placebo of -5.4 mmHg and -6.6 mmHg, respectively.

Table 17: Results from Placebo-Controlled Clinical Study of Canagliflozin as Add-on Therapy with Metformin¹

Efficacy Parameter	Canagliflozin + Metformin 26 weeks		Placebo + Metformin (N=183)
	100 mg (N=368)	300 mg (N=367)	
HbA1c (%)			
Baseline (mean)	7.94	7.95	7.96
Change from baseline (adjusted mean)	-0.79	-0.94	-0.17
Difference from placebo (adjusted mean) (95% CI)	-0.62 ² (-0.76; -0.48)	-0.77 ² (-0.91; -0.64)	N/A ³
Percent of patients achieving HbA1c < 7%	45.5 ²	57.8 ²	29.8
Fasting Plasma Glucose (mmol/L)			
Baseline (mean)	9.36	9.59	9.12
Change from baseline (adjusted mean)	-1.52	-2.10	0.14
Difference from placebo (adjusted mean) (95% CI)	-1.65 ² (-1.99; -1.32)	-2.23 ² (-2.57; -1.90)	N/A ³
2-hour Postprandial Glucose (mmol/L)			
Baseline (mean)	14.30	14.54	13.81
Change from baseline (adjusted mean)	-2.66	-3.17	-0.55
Difference from placebo (adjusted mean) (95% CI)	-2.12 ² (-2.73; -1.51)	-2.62 ² (-3.24; -2.01)	N/A ³
Body Weight			
Baseline (mean) in kg	88.7	85.4	86.7
% change from baseline (adjusted mean)	-3.7	-4.2	-1.2
Difference from placebo (adjusted mean) (95% CI)	-2.5 ² (-3.1; -1.9)	-2.9 ² (-3.5; -2.3)	N/A ³

¹ Intent-to-treat population using last observation in study prior to glycemic rescue therapy

² p<0.001 compared to placebo

³ N/A = Not applicable

Active-Controlled Study versus Glimpiride as add-on therapy with Metformin (Study DIA3009)

A total of 1450 patients with inadequate glycemic control (HbA1c level of $\geq 7\%$ to $\leq 9.5\%$) on metformin monotherapy ($\geq 2,000$ mg/day or at least 1,500 mg/day if higher dose not tolerated) participated in a randomized, double-blind, active-controlled, parallel-group, 3-arm, multicentre clinical study to evaluate the efficacy of canagliflozin as add-on therapy with metformin over 52 weeks. The mean age was 56 years, 52% of patients were men, and the mean baseline eGFR was 90 mL/min/1.73 m². Patients on metformin (N=928) at a stable protocol-specified dose entered a 2-week, single-blind, placebo run-in period. Other patients (N=522) entered a metformin dose titration and dose stabilization/antihyperglycemic agent washout period, immediately followed by the 2-week run-in period. Following the run-in period, patients with inadequate glycemic control were randomized to the addition of canagliflozin 100 mg, canagliflozin 300 mg, or glimepiride (titration allowed throughout the 52-week study to 6 to 8 mg), administered once daily.

As shown in Table 18 and Figure 4, after 52 weeks, treatment with canagliflozin 100 mg provided similar reductions in HbA1c from baseline compared to glimepiride (with the upper bound of the 95% confidence interval around the between-group difference less than the pre-specified non-inferiority margin of 0.3%); canagliflozin 300 mg provided a superior (p<0.05) reduction from baseline in HbA1c compared to glimepiride (with the upper bound of the 95%

confidence interval below 0). Statistically significant ($p < 0.001$) reductions in body weight were observed with canagliflozin compared to glimepiride. Reductions in systolic blood pressure were observed with canagliflozin 100 mg and 300 mg relative to glimepiride of -3.5 mmHg and -4.8 mmHg, respectively. The incidence of hypoglycemia with canagliflozin was significantly lower ($p < 0.001$) compared to glimepiride.

Table 18: Results from 52-Week Clinical Study Comparing Canagliflozin to Glimepiride as Add-on Therapy with Metformin¹

Efficacy Parameter	Canagliflozin + Metformin 52 Weeks		Glimepiride (titrated) + Metformin (N=482)
	100 mg (N=483)	300 mg (N=485)	
HbA1c (%)			
Baseline (mean)	7.78	7.79	7.83
Change from baseline (adjusted mean)	-0.82	-0.93	-0.81
Difference from glimepiride (adjusted mean) (95% CI)	-0.01 ² (-0.11; 0.09)	-0.12 ² (-0.22; -0.02)	N/A ³
Percent of patients achieving HbA1c < 7%	53.6	60.1	55.8
Fasting Plasma Glucose (mmol/L)			
Baseline (mean)	9.18	9.09	9.20
Change from baseline (adjusted mean)	-1.35	-1.52	-1.02
Difference from glimepiride (adjusted mean) (95% CI)	-0.33 (-0.56; -0.11)	-0.51 (-0.73; -0.28)	N/A ³
Body Weight			
Baseline (mean) in kg	86.8	86.6	86.6
% change from baseline (adjusted mean)	-4.2	-4.7	1.0
Difference from glimepiride (adjusted mean) (95% CI)	-5.2 ⁴ (-5.7; -4.7)	-5.7 ⁴ (-6.2; -5.1)	N/A ³


¹ Intent-to-treat population using last observation in study prior to glycemic rescue therapy

² Met pre-specified criteria for non-inferiority to glimepiride (with the upper bound of the 95% CI around the between-group difference less than the pre-specified non-inferiority margin of $< 0.3\%$). In a pre-specified assessment, the upper bound of the 95% CI for canagliflozin 300 mg, but not for canagliflozin 100 mg was < 0 , indicating a superior ($p < 0.05$) reduction in HbA1c relative to glimepiride with canagliflozin 300 mg.


³ N/A = Not applicable

⁴ $p < 0.001$

⁵ Includes only patients who had both baseline and post-baseline values



Note: LS Mean and SE in each post baseline visit are based on data with LOCF.



Note: LS Mean and SE in each post baseline visit are based on data with LOCF.

Figure 4: Mean Changes from Baseline for HbA1c (%) and Body Weight Over 52 Weeks in a Study Comparing Canagliflozin to Glimepiride as Add-on Therapy with Metformin

Add-on Therapy with Metformin and Sulfonylurea (Study DIA3002)

A total of 469 patients with inadequate glycemic control (HbA1c level of $\geq 7\%$ to $\leq 10.5\%$) on the combination of metformin (2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonylurea (maximal or near-maximal effective dose) participated in a randomized, double-blind, placebo-controlled, parallel-group, 3-arm, multicentre clinical study to evaluate the efficacy of canagliflozin as add-on therapy with metformin and sulfonylurea over 26 weeks. The mean age was 57 years, 51% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m². Patients on near-maximal or maximal effective doses of metformin and sulfonylurea (N=372) entered a 2-week, single-blind, placebo run-in period. Other patients

(N=97) entered a metformin and sulfonylurea dose titration and dose stabilization/antihyperglycemic agent washout period of up to 12 weeks, immediately followed by the 2-week run-in period. Following the run-in period, patients with inadequate glycemic control were randomized to the addition of canagliflozin 100 mg, canagliflozin 300 mg, or placebo administered once daily.

As shown in Table 19, statistically significant ($p < 0.001$) reductions in HbA1c, FPG, and body weight relative to placebo were observed. In addition, a greater percentage of patients achieved an HbA1c $< 7.0\%$ compared to placebo. Reductions in systolic blood pressure were observed with canagliflozin 100 mg and 300 mg relative to placebo of -2.2 mmHg and -1.6 mmHg, respectively. An increased incidence of hypoglycemia was observed in this study (see [7 WARNINGS AND PRECAUTIONS](#) and [8 ADVERSE REACTIONS](#)).

Table 19: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin as Add-on Therapy with Metformin and Sulfonylurea¹

Efficacy Parameter	Canagliflozin + Metformin and Sulfonylurea 26 Weeks		Placebo + Metformin and Sulfonylurea (N=156)
	100 mg (N=157)	300 mg (N=156)	
HbA1c (%)			
Baseline (mean)	8.13	8.13	8.12
Change from baseline (adjusted mean)	-0.85	-1.06	-0.13
Difference from placebo (adjusted mean) (95% CI)	-0.71 ² (-0.90; -0.52)	-0.92 ² (-1.11; -0.73)	N/A ³
Percent of patients achieving HbA1c < 7%	43.2 ²	56.6 ²	18.0
Fasting Plasma Glucose (mmol/L)			
Baseline (mean)	9.60	9.34	9.42
Change from baseline (adjusted mean)	-1.01	-1.69	0.23
Difference from placebo (adjusted mean) (95% CI)	-1.24 ² (-1.75; -0.73)	-1.92 ² (-2.43; -1.41)	N/A ³
Body Weight			
Baseline (mean) in kg	93.5	93.5	90.8
% change from baseline (adjusted mean)	-2.1	-2.6	-0.7
Difference from placebo (adjusted mean) (95% CI)	-1.4 ² (-2.1; -0.7)	-2.0 ² (-2.7; -1.3)	N/A ³

¹ Intent-to-treat population using last observation in study prior to glycemic rescue therapy

² $p < 0.001$ compared to placebo

³ N/A = Not applicable or not measured in this study

Active-Controlled Study versus Sitagliptin as Add-on Therapy with Metformin and Sulfonylurea (Study DIA3015)

A total of 755 patients with inadequate glycemic control (HbA1c level of $\geq 7.0\%$ to $\leq 10.5\%$) on the combination of metformin (2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonylurea (near-maximal or maximal effective dose) participated in a double-blind, active-controlled, parallel-group, 2-arm, multicentre clinical study to evaluate the efficacy of canagliflozin 300 mg as add-on therapy with metformin and sulfonylurea versus sitagliptin 100 mg as add-on therapy with metformin and sulfonylurea over 52 weeks. The mean age was 57 years, 56% of patients were men, and the mean baseline eGFR was 88 mL/min/1.73 m². Patients on near-maximal or maximal effective doses of metformin and sulfonylurea (N=716)

entered a 2-week single-blind, placebo run-in period. Other patients (N=39) entered a metformin and sulfonylurea dose titration and dose stabilization period of up to 12 weeks, immediately followed by the 2-week run-in period. Following the run-in period, patients with inadequate glycemic control were randomized to the addition of canagliflozin 300 mg or sitagliptin 100 mg.

As shown in Table 20 and Figure 5 after 52 weeks, canagliflozin 300 mg provided a superior ($p < 0.05$) reduction in HbA1c compared to sitagliptin 100 mg (with the upper bound of the 95% confidence interval around the between-group difference below 0). In addition, a greater percent of patients achieved an HbA1c of $< 7.0\%$ with canagliflozin 300 mg relative to sitagliptin: 47.6% of patients receiving canagliflozin 300 mg and 35.3% of patients receiving sitagliptin. Patients treated with canagliflozin 300 mg exhibited a significant mean decrease in percent change from baseline body weight compared to patients administered sitagliptin 100 mg. A statistically significant ($p < 0.001$) reduction in systolic blood pressure was observed with canagliflozin 300 mg of -5.9 mmHg relative to sitagliptin. A similar increased incidence of hypoglycemia was observed with both canagliflozin 300 mg and sitagliptin in this study, consistent with the expected increase of hypoglycemia when agents not associated with hypoglycemia are added to sulfonylurea (see [7 WARNINGS AND PRECAUTIONS](#) and [8 ADVERSE REACTIONS](#)). The proportion of patients who met glycemic withdrawal criteria (based on FPG until Week 26 and HbA1c thereafter) was lower with canagliflozin 300 mg (10.6%) compared with sitagliptin 100 mg (22.5%).

Table 20: Results from 52-Week Clinical Study Comparing Canagliflozin to Sitagliptin as Add-on Therapy with Metformin and Sulfonyleurea¹


Efficacy Parameter	Canagliflozin 300 mg + Metformin and Sulfonyleurea (N=377)	Sitagliptin 100 mg + Metformin and Sulfonyleurea (N=378)
HbA1c (%)		
Baseline (mean)	8.12	8.13
Change from baseline (adjusted mean)	-1.03	-0.66
Difference from sitagliptin (adjusted mean) (95% CI)	-0.37 ² (-0.50; -0.25)	N/A ⁴
Percent of patients achieving HbA1c < 7%	47.6	35.3
Fasting Plasma Glucose (mmol/L)		
Baseline (mean)	9.42	9.09
Change from baseline (adjusted mean)	-1.66	-0.32
Difference from sitagliptin (adjusted mean) (95% CI)	-1.34 (-1.66; -1.01)	N/A ⁴
Body Weight		
Baseline (mean) in kg	87.6	89.6
% change from baseline (adjusted mean)	-2.5	0.3
Difference from sitagliptin (adjusted mean) (95% CI)	-2.8 ³ (-3.3; -2.2)	N/A ⁴

¹ Intent-to-treat population using last observation in study prior to glycemic rescue therapy

² Met pre-specified criteria for non-inferiority to sitagliptin (with the upper bound of the 95% CI around the between-group difference less than the pre-specified non-inferiority margin of < 0.3%); in a pre-specified assessment, the upper bound of the 95% CI for canagliflozin 300 mg was < 0, indicating a superior (p<0.05) reduction in HbA1c relative to sitagliptin with canagliflozin 300 mg.

³ p<0.001

⁴ N/A = Not applicable



Note: LS Mean and SE in each post-baseline visit are based on data with LOCF.

Figure 5: Mean Change from Baseline for HbA1c (%) Over 52 Weeks in a Study Comparing Canagliflozin to Sitagliptin as Add-on Therapy with Metformin and Sulfonyleurea

Add-on Therapy with Metformin and Pioglitazone (Study DIA3012)

A total of 342 patients with inadequate glycemic control (HbA1c level of $\geq 7.0\%$ to $\leq 10.5\%$) on the combination of metformin (2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and pioglitazone (30 or 45 mg/day) participated in a randomized, double-blind, placebo-controlled, parallel-group, 3-arm, multicentre clinical study to evaluate the efficacy of canagliflozin as add-on therapy with metformin and pioglitazone over 26 weeks. The mean age was 57 years, 63% of patients were men, and the mean baseline eGFR was 86 mL/min/1.73 m². Patients already on protocol-specified doses of metformin and pioglitazone (N=163) entered a 2-week, single-blind, placebo run-in period. Other patients (N=181) entered a metformin and pioglitazone dose titration and dose stabilization period for up to 12 weeks with at least 8 weeks on stable doses of metformin and pioglitazone, immediately followed by the 2-week run-in period. Following the run-in period, patients with inadequate glycemic control were randomized (N=344) to the addition of canagliflozin 100 mg, canagliflozin 300 mg, or placebo, administered once daily.

As shown in Table 21, statistically significant ($p < 0.001$) reductions in HbA1c, baseline FPG, and body weight relative to placebo were observed for canagliflozin at Week 26. In addition, a greater percent of patients achieved an HbA1c of $< 7.0\%$ compared to placebo. Statistically significant reductions in systolic blood pressure were observed with canagliflozin 100 mg and 300 mg relative to placebo of -4.1 mmHg ($p = 0.005$) and -3.5 mmHg ($p = 0.016$), respectively.

Table 21: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin as Add-on Therapy with Metformin and Pioglitazone¹

Efficacy Parameter	Canagliflozin + Metformin and Pioglitazone 26 Weeks		Placebo + Metformin and Pioglitazone (N=115)
	100 mg (N=113)	300 mg (N=114)	
HbA1c (%)			
Baseline (mean)	7.99	7.84	8.00
Change from baseline (adjusted mean)	-0.89	-1.03	-0.26
Difference from placebo (adjusted mean) (95% CI)	-0.62 ² (-0.81; -0.44)	-0.76 ² (-0.95; -0.58)	N/A ³
Percent of patients achieving HbA1c < 7%	46.9²	64.3²	32.5
Fasting Plasma Glucose (mmol/L)			
Baseline (mean)	9.38	9.11	9.13
Change from baseline (adjusted mean)	-1.49	-1.84	0.14
Difference from placebo (adjusted mean) (95% CI)	-1.63 ² (-2.05; -1.21)	-1.98 ² (-2.41; -1.56)	N/A ³
Body Weight			
Baseline (mean) in kg	94.2	94.4	94
% change from baseline (adjusted mean)	-2.8	-3.8	-0.1
Difference from placebo (adjusted mean) (95% CI)	-2.7 ² (-3.6; -1.8)	-3.7 ² (-4.6; -2.8)	N/A ³

¹ Intent-to-treat population using last observation in study prior to glycemic rescue therapy

² p<0.001 compared to placebo

³ N/A = Not applicable or not measured in this study

Add-on Therapy with Insulin and Metformin (Derived from DIA3008 substudy)

A total of 1718 patients with inadequate glycemic control (HbA1c level of ≥ 7.0 to $\leq 10.5\%$) on insulin ≥ 30 units/day or insulin add-on therapy with other antihyperglycemic agents participated in a randomized, double-blind, placebo-controlled, parallel-group, 3-arm, multicentre substudy of a cardiovascular outcomes study; this substudy evaluated the efficacy of canagliflozin as add-on therapy with insulin over 18 weeks. The mean age was 63 years, 66% of patients were men, and the mean baseline eGFR was 75 mL/min/1.73 m². Patients on basal, bolus, or basal/bolus insulin, with the majority on a background basal/bolus insulin regimen, for at least 10 weeks entered a 2-week, single-blind, placebo run-in period. After the run-in period, patients with inadequate glycemic control were randomized to the addition of canagliflozin 100 mg, canagliflozin 300 mg, or placebo, administered once daily. The mean daily insulin dose at baseline was 83 units, which was similar across treatment groups.

A subgroup of 731 patients with inadequate glycemic control received canagliflozin in combination with metformin and ≥ 30 units/day of insulin over 18 weeks. As shown in Table 22, statistically significant (p<0.001) reductions in HbA1c, FPG, and body weight relative to placebo were observed for canagliflozin at Week 18 in patients on an insulin+metformin background. In addition, a greater percentage of patients achieved an HbA1c < 7.0% compared to placebo. Reductions in systolic blood pressure were observed with canagliflozin 100 mg and 300 mg relative to placebo of -2.9 mmHg (p=0.011) and -4.8 mmHg (p<0.001), respectively. An increased incidence of hypoglycemia was observed in this study (see [7 WARNINGS AND PRECAUTIONS](#), [8 ADVERSE REACTIONS](#), and [4 DOSAGE AND ADMINISTRATION](#)).

Table 22: Results from 18-Week Placebo-Controlled Clinical Study of Canagliflozin as Add-on Therapy with Insulin ≥ 30 Units/Day (With Insulin and Metformin)¹

Efficacy Parameter	Canagliflozin + Insulin + Metformin 18 Weeks		Placebo + Insulin + Metformin (N=244)
	100 mg (N=241)	300 mg (N=246)	
HbA1c (%)			
Baseline (mean)	8.28	8.21	8.21
Change from baseline (adjusted mean)	-0.66	-0.77	0.01
Difference from placebo (adjusted mean) (95% CI)	-0.67 ² (-0.79; -0.55)	-0.78 ² (-0.90; -0.66)	N/A ³
Percent of patients achieving HbA1c < 7%	19.6 ²	26.7 ²	7.1
Fasting Plasma Glucose (mmol/L)			
Baseline	9.38	9.35	9.34
Change from baseline (adjusted mean)	-1.06	-1.48	0.09
Difference from placebo (adjusted mean) (95% CI)	-1.15 ² (-1.56; -0.73)	-1.57 ² (-1.98; -1.16)	N/A ³
Body Weight			
Baseline (mean) in kg	97.4	98.4	99.9
% change from baseline (adjusted mean)	-1.9	-2.7	0.0
Difference from placebo (adjusted mean) (95% CI)	-1.9 ² (-2.4; -1.5)	-2.7 ² (-3.2; -2.3)	N/A ³

¹ Intent-to-treat population using last observation in study prior to glycemic rescue therapy

² p < 0.001 compared to placebo

³ N/A = Not applicable

Cardiovascular Outcomes (CANVAS (DIA3008) and CANVAS-R (DIA4003))

The effect of canagliflozin on cardiovascular risk in adults with type 2 diabetes who had established cardiovascular (CV) disease or were at risk for CVD (two or more CV risk factors), was evaluated in the CANVAS Program (CANVAS and CANVAS-R studies). These studies were multicenter, multi-national, randomized, double-blind, placebo-controlled parallel group, time- and event-driven, with similar inclusion and exclusion criteria and patient populations. The studies compared the risk of experiencing a Major Adverse Cardiovascular Event (MACE) defined as the composite of cardiovascular death, nonfatal myocardial infarction and nonfatal stroke, between canagliflozin and placebo on a background of standard of care treatments for diabetes and atherosclerotic cardiovascular disease. Additional pre-specified, adjudicated endpoints included CV death, fatal/non-fatal myocardial infarction, fatal/non-fatal stroke, hospitalization for heart failure, and all-cause mortality.


In CANVAS, subjects were randomly assigned 1:1:1 to canagliflozin 100 mg, canagliflozin 300 mg, or matching placebo. In CANVAS-R, subjects were randomly assigned 1:1 to canagliflozin 100 mg or matching placebo, and titration to 300 mg was permitted at the investigator's discretion (based on tolerability and glycemic needs) at Week 13 or later visits. Concomitant antidiabetic and atherosclerotic therapies could be adjusted, at the discretion of investigators, to ensure participants were treated according to the standard care for these diseases.

A total of 10,134 patients were treated (4,327 in CANVAS and 5,807 in CANVAS-R; total of 4,344 randomly assigned to placebo and 5,790 to canagliflozin). For the integrated CANVAS

trials, the mean duration of treatment was 149.2 weeks (mean of 222.8 weeks for CANVAS and 94.4 weeks for CANVAS-R) and the mean duration of study follow-up was 188.2 weeks (mean of 295.9 for CANVAS and 108.0 weeks for CANVAS-R). Vital status was obtained for 99.6% of the subjects. The proportion of subjects who completed the study was 96.0%. Approximately 78% of the study population was Caucasian, 13% was Asian, and 3% was Black. The mean age was 63 years and approximately 64% were male. All patients in the study had inadequately controlled type 2 diabetes mellitus at baseline ($\text{HbA}_{1c} \geq 7.0\%$ to $\leq 10.5\%$). The mean HbA_{1c} at baseline was 8.2% and mean duration of diabetes was 13.5 years. Baseline renal function was normal or mildly impaired in 80% of patients and moderately impaired in 20% of patients (mean eGFR 77 mL/min/1.73 m²). There were 526 patients with eGFR 30-<45 mL/min/1.73 m², 1485 patients with eGFR 45-<60 mL/min/1.73 m², and 5625 with eGFR 60-<90 mL/min/1.73 m². At baseline, 99% of patients were treated with one or more antidiabetic medications including metformin (77%), insulin (50%), and sulfonylurea (43%).

Sixty-six percent of subjects had a history of established cardiovascular disease, with 56% having a history of coronary disease, 19% with cerebrovascular disease, and 21% with peripheral vascular disease; 14% had a history of heart failure. At baseline, the mean systolic blood pressure was 137 mmHg, the mean diastolic blood pressure was 78 mmHg, the mean LDL was 2.29 mmol/L, the mean HDL was 1.2 mmol/L, and the mean urinary albumin to creatinine ratio (UACR) was 115 mg/g. At baseline, approximately 80% of patients were treated with renin angiotensin system inhibitors, 54% with beta-blockers, 13% with loop diuretics, 36% with non-loop diuretics, 75% with statins, and 74% with antiplatelet agents (including aspirin).

The primary endpoint in the CANVAS Program was the time to first occurrence of a composite MACE endpoint of cardiovascular death, nonfatal myocardial infarction, or nonfatal stroke, considering all events up to individual trial completion. The MACE hazard ratio (HR) in patients treated with canagliflozin compared with placebo and its 95% CI was estimated using a stratified Cox proportional hazards regression model with stratification by study and by established cardiovascular disease (HR: 0.86; 95% CI 0.75, 0.97, $p < 0.0001$ for non-inferiority; $p = 0.0158$ for superiority). According to the primary hypothesis, the integrated canagliflozin treatment (CANVAS and CANVAS-R) was found to be non-inferior to placebo, since the upper bound of the 95% CI was below 1.3 and superior to placebo, since the upper bound of the 95% CI was also below 1.0. Each of the components of the MACE composite endpoint showed a similar reduction when assessed as independent endpoints (see Figure 6). Results for the 100 mg and 300 mg canagliflozin doses were consistent with results for the combined dose groups. The reduction in MACE was accounted for by the subgroup of patients with established cardiovascular disease (HR 0.82; 95% CI 0.72, 0.95) (see Figure 6), whilst the subgroup of patients with only risk factors for cardiovascular disease at baseline had a hazard ratio whose 95% confidence interval included one (HR 0.98; 95% CI 0.74, 1.30).



¹ P value for superiority (2-sided) = 0.0158.

Figure 6: Treatment Effect for the Primary Composite Endpoint and its Components

Based on the Kaplan-Meier plot for the first occurrence of MACE, shown below, the reduction in MACE in the canagliflozin group was observed as early as Week 26 and was maintained throughout the remainder of the study (Figure 7 and Figure 8)




Figure 7: Time to First Occurrence of MACE (CANVAS Integrated)




Figure 8: Time to First Occurrence of MACE (Subjects with Established CV Disease)

In the CANVAS program, subjects treated with canagliflozin had a lower risk of hospitalization for heart failure compared to those treated with placebo.

Table 23: Treatment Effect for Hospitalized Heart Failure and the Composite of Cardiovascular Death or Hospitalized Heart Failure

	Placebo N=4347 Event rate per 100 patient- years	Canagliflozin N=5795 Event rate per 100 patient- years	Hazard ratio vs. Placebo (95% CI)
Hospitalized heart failure (time to first occurrence; intent-to-treat analysis set)	0.87	0.55	0.67 (0.52, 0.87) ¹
Death or Hospitalization due to heart failure (time to first occurrence; intent-to-treat analysis set)	0.97	0.64	0.70 (0.55, 0.89)

¹ p=0.0021; nominal value




Figure 9: Time to First Occurrence of Hospitalization of Heart Failure

Renal and Cardiovascular Outcomes in Patients with Type 2 Diabetes Mellitus and Diabetic Nephropathy (CREDESCENCE DNE3001)

The Canagliflozin and Renal Events in Diabetes with Established Nephropathy Clinical Evaluation Trial (CREDESCENCE) studied the effect of canagliflozin 100 mg relative to placebo on progression to end-stage kidney disease (ESKD), doubling of serum creatinine, and renal or cardiovascular (CV) death in adults with type 2 diabetes and diabetic nephropathy with (eGFR) ≥ 30 to < 90 mL/min/1.73 m² and albuminuria (> 33.9 to ≤ 565.6 mg/mmol of creatinine), who were receiving standard of care including maximally tolerated labelled dose of an angiotensin-converting enzyme inhibitor (ACEi) or angiotensin receptor blocker (ARB). This study was a multicenter, randomized, double-blind, event-driven, placebo-controlled, parallel-group, 2-arm study.

In CREDESCENCE, subjects were randomly assigned 1:1 to canagliflozin 100mg or placebo, stratified by screening estimated glomerular filtration rate (eGFR) ≥ 30 to < 45 , ≥ 45 to < 60 , ≥ 60 to < 90 mL/min/1.73 m². Treatment with canagliflozin 100 mg was continued in patients until the initiation of dialysis or renal transplantation.

A total of 4,401 subjects were randomized (2,199 randomly assigned to placebo and 2,202 to canagliflozin 100mg), followed for a mean duration of 136 weeks, and included in the intent-to-treat analysis set. Four of the randomized subjects were not dosed, leading to 4,397 subjects (exposed for a mean duration of 115 weeks) in the on-treatment analysis set. Vital status was obtained for 99.9% of subjects across the study. The majority (67%) of the study population identified as White, 20% as Asian, and 5% as Black; 32% of all subjects were of Hispanic or Latino ethnicity. The mean age was 63 years and approximately 66% were male.

The mean baseline HbA1c was 8.3%, with 53.2% of subjects having baseline HbA1c $\geq 8\%$, and the baseline median urine albumin/creatinine was 104.75 mg/mmol. The most frequent antihyperglycemic agents (AHA) medications used at baseline were insulin (65.5%), biguanides (57.8%), and sulfonylureas (28.8%). Nearly all subjects (99.9%) were on ACEi or ARB at randomization. About 92% of the subjects were on cardiovascular therapies (not including ACEi/ARBs) at baseline, with approximately 60% taking an anti-thrombotic agent (including aspirin) and 69% on statins.


The mean baseline eGFR was 56.2 mL/min/1.73 m² and approximately 60% of the population had a baseline eGFR of < 60 mL/min/1.73 m². Subjects had a mean duration of diabetes of approximately 16 years. The proportion of subjects with prior CV disease was 50.4%; 14.8% had a history of heart failure. While the entire study population had nephropathy at baseline, about 64% of the population had at least 2 microvascular complications (i.e. diabetic nephropathy and another microvascular complication). At baseline, 5.4% of subjects in the canagliflozin 100mg arm had a history of amputation and 5.2% of subjects in the placebo arm.

The primary composite endpoint in the CREDENCE study was the time to first occurrence of ESKD (defined as an eGFR < 15 mL/min/1.73 m², initiation of chronic dialysis or renal transplant), doubling of serum creatinine, and renal or CV death. Canagliflozin 100 mg significantly reduced the risk of first occurrence of the primary composite endpoint of ESKD, doubling of serum creatinine, and renal or CV death [$p < 0.0001$; HR:0.70; 95% CI:0.59, 0.82] (see Figure 10 and Figure 11). The treatment effect reflected a reduction in progression to ESKD, doubling of serum creatinine and cardiovascular death. There were few renal deaths during the trial. The efficacy of canagliflozin 100 mg on the primary endpoint composite was generally consistent across major demographic and disease subgroups, including a subgroup defined by the 3 screening eGFR strata.

Canagliflozin 100 mg significantly reduced the risk of the following secondary endpoints, as shown in Figure 10 below: Composite endpoint of CV Death and Hospitalized Heart Failure [HR:0.69; 95% CI: 0.57 to 0.83; $p = 0.0001$], MACE (Major Adverse Cardiovascular Events) (comprised of non-fatal MI, non-fatal stroke and CV death) [HR:0.80; 95% CI:0.67 to 0.95; $p = 0.0121$], Hospitalized Heart Failure [HR:0.61; 95% CI:0.47 to 0.80; $p = 0.0003$], and Renal composite endpoint (comprised of ESKD, doubling of serum creatinine, and renal death) [HR:0.66; 95% CI:0.53 to 0.81; $p < 0.0001$].

For both primary and secondary endpoints, the HR in subjects treated with canagliflozin 100 mg compared with placebo and its 95% CI were estimated using a stratified Cox proportional hazards regression model with treatment as the explanatory variable and stratified by screening eGFR (≥ 30 to < 45 , ≥ 45 to < 60 , ≥ 60 to < 90 mL/min/1.73 m²).

Forest Plot of Hazard Ratios and 95% CI of the Primary Composite Endpoint, Secondary Endpoints, and Their Components (Intention-to-Treat Analysis Set)



CI, confidence interval; ESKD, end-stage kidney disease; CV, cardiovascular; NS, not significant; HHF, hospitalization for heart failure; MI, myocardial infarction. MACE is the 3-point Major Adverse Cardiac Event (CV death, nonfatal MI, and nonfatal stroke).

The individual components do not represent a breakdown of the composite outcomes, but rather the total number of subjects experiencing an event during the course of the study.

*Testing of the primary and the secondary efficacy endpoints was performed using a 2-sided alpha level of 0.022 and 0.038, respectively.

[†]CV death is being presented as a component of the primary composite endpoint, as a component of MACE, and as a secondary endpoint which underwent formal hypothesis testing.

[‡]Fatal/nonfatal MI and fatal/nonfatal stroke were not prespecified in the hierarchical testing sequence and are considered exploratory endpoints.

Figure 10: Treatment Effect for the Primary and Secondary Composite Endpoints and their Components

Based on the Kaplan-Meier plot for the time to first occurrence of the primary composite endpoint of ESKD, doubling of serum creatinine, renal death, and CV death shown below, the curves began to separate by Week 52 and continued to diverge thereafter (see Figure 11).




Figure 11: Time to First Occurrence of the Primary Composite Endpoint (ESKD, Doubling of Serum Creatinine, Renal Death, CV Death)

The Kaplan-Meier plot for the first occurrence of hospitalized heart failure over time is shown in Figure 12. Canagliflozin significantly reduced the risk of hospitalized heart failure as compared with placebo (HR: 0.61; 95% CI: 0.47, 0.80; p=0.0003). The Kaplan-Meier curves separated within the first 26 weeks of treatment and continued to diverge thereafter.




Figure 12: Time to First Occurrence of Hospitalized Heart Failure

The Kaplan-Meier plot for the first occurrence of the secondary renal composite endpoint of doubling of serum creatinine, ESKD, and renal death over time is shown in Figure 13. Canagliflozin significantly reduced the risk of the secondary renal composite endpoint as compared with placebo (HR: 0.66; 95% CI: 0.53, 0.81; $p < 0.0001$). The Kaplan-Meier curves separated within the first 52 weeks of treatment and continued to diverge thereafter.




Figure 13: Time to First Occurrence of Renal Composite Endpoint (Doubling of Serum Creatinine/ESKD/Renal Death)

15 MICROBIOLOGY

No microbiological information is required for this product.

16 NON-CLINICAL TOXICOLOGY

General Toxicology

Canagliflozin

Non-clinical data reveal no particular hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, and genotoxicity.

Canagliflozin has relatively low acute oral toxicity, with maximum non-lethal single doses of 2000 mg/kg in mice (both sexes) and male rats, and 1000 mg/kg in female rats.

Repeat-dose oral toxicity studies were conducted in mice, rats and dogs for up to 3, 6 and 12 months, respectively. Canagliflozin was generally well tolerated up to oral doses of 4 mg/kg/day in rats and 100 mg/kg/day in mice and dogs (up to approximately 0.5, 11, and 20 times the clinical dose of 300 mg based on AUC exposure for rats, mice and dogs, respectively). The major adverse effects, observed mainly in rats, were related to the pharmacologic mode of action of canagliflozin, and these included increased urinary glucose, increased urine volume, increased urinary excretion of electrolytes, decreased plasma glucose at high dose levels, and reduced body weight. The primary targets of toxicity were the kidney

and bone. In the 3-month rat study, minimal mineralization of renal interstitium and/or pelvis were observed in some animals given doses of ≥ 4 mg/kg/day. In the 6-month rat study, renal tubular dilatation was seen at all doses (4, 20 and 100 mg/kg/day), and an increased incidence and severity of transitional epithelial hyperplasia in the renal pelvis was observed at 100 mg/kg/day. In dogs, treatment-related tubular regeneration/degeneration and tubular dilatation occurred only at the high dose of 200/100 mg/kg/day. Trabecular hyperostosis was observed in the repeat-dose studies in rats, but not in mice and dogs. In the 2-week rat study, canagliflozin at 150 mg/kg/day caused minimal to mild hyperostosis but in 3- and 6-month rat studies, hyperostosis was detected at 4 mg/kg/day, the lowest dose tested. A 1-month mechanistic rat study showed that hyperostosis occurred in young, actively growing animals (6 to 8 weeks old, as in the toxicity studies) but not in older (6 month old) animals where bone growth has substantially slowed.

Carcinogenicity

Canagliflozin and Metformin hydrochloride Combination

No animal studies have been conducted with the combined products in INVOKAMET to evaluate carcinogenesis.

Canagliflozin

The carcinogenicity of canagliflozin was evaluated in 2-year studies in mice and rats at oral doses of 10, 30, or 100 mg/kg/day. Canagliflozin did not increase the incidence of tumors in male and female mice up to 100 mg/kg/day (up to 14 times the clinical dose of 300 mg based on AUC exposure).

The incidence of testicular Leydig cell tumors increased significantly in male rats at all doses tested (≥ 1.5 times the clinical dose of 300 mg based on AUC exposure). The Leydig cell tumors are associated with an increase in luteinizing hormone (LH), which is a known mechanism of Leydig cell tumor formation in rats. In a 12-week clinical study, unstimulated LH did not increase in males treated with canagliflozin.

The incidence of pheochromocytomas and renal tubular tumors increased significantly in male and female rats given high doses of 100 mg/kg/day (approximately 12 times the clinical dose of 300 mg based on AUC exposure). Canagliflozin-induced renal tubule tumors and pheochromocytomas in rats may be caused by carbohydrate malabsorption; mechanistic clinical studies have not demonstrated carbohydrate malabsorption in humans at canagliflozin doses of up to 2 times the recommended clinical dose of 300 mg.

Metformin hydrochloride

Long-term carcinogenicity studies have been performed in rats (dosing duration of 104 weeks) and mice (dosing duration of 91 weeks) at doses up to and including 900 mg/kg/day and 1500 mg/kg/day, respectively. These doses are both approximately 4 times the maximum recommended human daily dose of 2000 mg based on body surface area comparisons. No evidence of carcinogenicity with metformin was found in either male or female mice. Similarly, there was no tumorigenic potential observed with metformin in male rats. There was, however, an increased incidence of benign stromal uterine polyps in female rats treated with 900 mg/kg/day.

Genotoxicity

Canagliflozin and Metformin hydrochloride Combination

No animal studies have been conducted with the combined products in INVOKAMET to evaluate mutagenesis.

Canagliflozin

Canagliflozin was not mutagenic with or without metabolic activation in the Ames assay. Canagliflozin was mutagenic in the *in vitro* mouse lymphoma assay with but not without metabolic activation. Canagliflozin was not mutagenic or clastogenic in an *in vivo* oral micronucleus assay in rats and an *in vivo* oral Comet assay in rats.

Metformin hydrochloride

There was no evidence of a mutagenic potential of metformin in the following *in vitro* tests: Ames test (*S. typhimurium*), gene mutation test (mouse lymphoma cells), or chromosomal aberrations test (human lymphocytes). Results in the *in vivo* mouse micronucleus test were also negative.

Reproductive and Developmental Toxicology

Canagliflozin and Metformin hydrochloride Combination

No animal studies have been conducted with the combined products in INVOKAMET to evaluate impairment of fertility.

Canagliflozin

In rat fertility studies, canagliflozin had no adverse effects on mating, fertility, or early embryonic development up to the highest dose of 100 mg/kg/day (up to 19 times the clinical dose of 300 mg based on AUC exposure), although there were slight sperm morphological changes at this dose level.

Canagliflozin was not teratogenic at any dose tested when administered orally to pregnant rats and rabbits during the period of organogenesis. In both rats and rabbits, a slight increase in the number of fetuses with reduced ossification, indicative of a slight developmental delay, was observed at the high doses (approximately 19 times the clinical dose of 300 mg based on AUC exposure) in the presence of maternal toxicity.

In a pre- and postnatal development study, canagliflozin administered orally to female rats from gestation Day 6 to lactation Day 20 resulted in decreased body weights in male and female offspring at maternally toxic doses of ≥ 30 mg/kg/day (≥ 5.9 times the clinical dose of 300 mg based on AUC exposure). Maternal toxicity was limited to decreased body weight gain.

Metformin hydrochloride

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately 3 times the maximum recommended human daily dose of 2000 mg based on body surface area comparisons.

Metformin was not teratogenic in rats and rabbits at doses up to 600 mg/kg/day. This represents an exposure of about 2 and 6 times the maximum recommended human daily dose of 2,000 mg based on body surface area comparisons for rats and rabbits, respectively. Determination of

fetal concentrations demonstrated a partial placental barrier to metformin.

Juvenile Toxicity

Canagliflozin

In a juvenile toxicity study in which canagliflozin was dosed orally to young rats from postnatal day (PND) 21 until PND 90 at doses of 4, 20, 65, or 100 mg/kg, increased kidney weights and a dose-related increase in the incidence and severity of renal pelvic and renal tubular dilatation were reported at all dose levels. Exposure at the lowest dose tested was approximately 0.5 times the maximum recommended clinical dose of 300 mg. The renal pelvic dilatations observed in juvenile animals did not fully reverse within the 1-month recovery period. Persistent renal findings in juvenile rats can most likely be attributed to reduced ability of the developing rat kidney to handle canagliflozin-increased urine volumes, as functional maturation of the rat kidney continues through 6 weeks of age. Additionally, shortened ulna growth and delays in sexual maturation were observed in juvenile rats at doses that were greater than or equal to 3 times and 9 times the clinical dose of 300 mg based on AUC exposure, respectively.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

P^rINVOKAMET[®] canagliflozin and metformin hydrochloride tablets

Read this carefully before you start taking **INVOKAMET** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **INVOKAMET**.

Serious Warnings and Precautions

Lactic Acidosis

- INVOKAMET contains the medicinal ingredient metformin. Having too much metformin in your blood can cause lactic acidosis. Lactic acidosis is a rare but serious build-up of lactic acid in the blood. It can cause death and must be treated in a hospital setting.
- Alcohol may increase your risk of lactic acidosis. Do not drink a lot of alcohol while taking INVOKAMET. This means that you should not “binge” drink (have 5 or more drinks in a row by men, or 4 or more drinks in a row by women).

Diabetic Ketoacidosis (DKA)

- DKA may happen during or after stopping treatment with INVOKAMET. It is a serious and life-threatening condition, which may need urgent hospital care. Some cases of DKA have led to death. DKA is a complication of diabetes, where your body produces high levels of blood acids called ketones. It can happen in patients with type 2 diabetes mellitus (T2DM), with normal or high blood sugar (glucose) levels, who are treated with INVOKAMET or with other sodium-glucose co-transporter 2 (SGLT2) inhibitors.
- **Seek medical attention right away and stop taking INVOKAMET immediately** if you have any of the following symptoms (even if your blood sugar levels are normal): difficulty breathing, nausea, vomiting, stomach pain, loss of appetite, confusion, feeling very thirsty, feeling unusually tired or sleepy, a sweet smell to the breath, a sweet or metallic taste in the mouth, or a different odour to urine or sweat.
- Do NOT use INVOKAMET if you have:
 - type 1 diabetes
 - DKA or a history of DKA

Lower Limb Amputation

- INVOKAMET may increase your risk of lower limb amputations. Amputations have happened mainly on the toe or part of the foot, however, amputations involving the leg, below and above the knee have also occurred. Some people had more than one amputation, some on both sides of the body.
- Tell your healthcare professional if you have ever had an amputation, blood vessel disease, nerve disease, or a foot ulcer (sore) caused by diabetes.

- **Seek medical attention right away** if you have new pain or tenderness, any sores, ulcers, or infections in your leg or foot. Your healthcare professional may decide to stop your INVOKAMET if you have any of these signs or symptoms. Talk to your healthcare professional about proper foot care and keeping hydrated.

What is INVOKAMET used for?

INVOKAMET is used along with diet and exercise to improve blood sugar levels in adults with type 2 diabetes.

INVOKAMET can be used in patients:

- whose diabetes is not controlled on:
 - metformin alone
 - metformin in combination with a sulfonylurea
 - metformin in combination with pioglitazone
 - metformin in combination with insulin
- who are currently taking combination of separate tablets of:
 - metformin and INVOKANA (canagliflozin)
 - metformin, INVOKANA and a sulfonylurea
 - metformin, INVOKANA and pioglitazone
 - metformin, INVOKANA and insulin

How does INVOKAMET work?

INVOKAMET contains two different medicines, canagliflozin and metformin. They work together to help you better control your blood sugar level.

Canagliflozin is in a class of medicines called sodium-glucose co-transporter 2 (SGLT2) inhibitors. It lowers blood sugar by causing the kidneys to remove more sugar in the urine.

Metformin is in a class of medicines called biguanides. It helps your body respond better to insulin, a natural chemical that manage the amount of sugar in the blood. This lowers the amount of sugar made by the liver and the amount of sugar that moves from the food you eat to the blood.

What are the ingredients in INVOKAMET?

Medicinal ingredients: canagliflozin and metformin hydrochloride

Non-medicinal ingredients: croscarmellose sodium, hypromellose, iron oxide black (150/1000 mg tablets), iron oxide red (50 mg/1000 mg, 150 mg/500 mg and 150 mg/1000 mg tablets), iron oxide yellow (50 mg/1000 mg, and 150 mg/500 mg), macrogol (polyethylene glycol), magnesium stearate, microcrystalline cellulose, polyvinyl alcohol (partially hydrolyzed), talc, and titanium dioxide

INVOKAMET comes in the following dosage forms:

Tablets:

- 50 mg canagliflozin / 500 mg metformin hydrochloride
- 50 mg canagliflozin / 1000 mg metformin hydrochloride
- 150 mg canagliflozin / 500 mg metformin hydrochloride

- 150 mg canagliflozin / 1000 mg metformin hydrochloride

Do not use INVOKAMET if you:

- are allergic to canagliflozin, metformin or any other ingredients in INVOKAMET.
- have severe kidney problems or end-stage kidney disease.
- are on dialysis.
- do not know the health status of your kidneys.
- have or have had a condition called metabolic acidosis (including diabetic ketoacidosis, or lactic acidosis – too much acid in the blood).
- have type 1 diabetes (your body does not produce any insulin).
- drink alcohol very often or drink a lot of alcohol in a short-term “binge” drinking.
- have liver problems.
- have serious heart problems or heart failure.
- have lost a lot of water from your body (seriously dehydrated).
- are in shock.
- are under stress, have a serious infection, have recently had an injury.
- are going to have major surgery or are recovering from surgery.
- are going to have an exam or other tests such as an X-ray or scan with injectable dyes or contrast agents used. You will need to stop taking this medicine around the time of your procedure.
- are pregnant or are planning to become pregnant. It is not known if INVOKAMET can harm your unborn baby.
- are breastfeeding.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take INVOKAMET. Talk about any health conditions or problems you may have, including if you:

- **have any of the following conditions:**
 - are malnourished.
 - problems with your adrenal or pituitary glands (adrenal or pituitary insufficiency).
 - low vitamin B₁₂.
 - history of yeast infections of the vagina or penis.
 - current or history of low blood pressure (hypotension).
- are taking any of the medications listed in the drug interactions section below (see **The following may interact with INVOKAMET**).
- **have risk factors for DKA, such as if you:**
 - are dehydrated or suffer from excessive vomiting, diarrhea, or sweating.
 - are on a very low carbohydrate (sugar) diet.
 - have been fasting for a while.
 - are eating less, or there is a change in your diet.
 - drink a lot of alcohol.
 - have or have had problems with your pancreas, including pancreatitis or surgery on your pancreas.
 - are hospitalized for major surgery, or are about to have major surgery.
 - are hospitalized for serious infection or serious medical illnesses.
 - need to have a procedure that requires long periods of fasting.

- have an acute illness.
 - have sudden reduction in insulin dose.
 - have diabetic kidney disease. This is when your kidneys are damaged as a result of your diabetes.
 - have a history of DKA.
- **have risk factors for amputation, such as if you:**
 - have a history of amputation.
 - have heart disease or are at risk for heart disease.
 - have had blocked or narrowed blood vessels, usually in your leg.
 - have damage to the nerves (neuropathy) in your leg.
 - have had diabetic foot ulcers or sores.
 - have a lower limb infection.
 - are dehydrated.
- **have risk factors for lactic acidosis, such as if you:**
 - have severe kidney problems.
 - have liver problems.
 - have congestive heart failure that requires treatment with medicines.
 - have a heart attack, severe infection, or stroke.
 - drink a lot of alcohol (very often or short-term “binge” drinking).
 - are dehydrated. Dehydration can also happen when you sweat a lot with activity or exercise and don’t drink enough fluids. Tell your healthcare professional if this happens.
 - have certain x-ray tests with injectable dyes or contrast agents used.
 - have surgery.

Other warnings you should know about:

Vitamin B₁₂ levels

INVOKAMET can cause your vitamin B₁₂ levels to be low. This can cause peripheral neuropathy (nerve damage) or anemia (low red blood cells).

Children and adolescents (under 18 years of age)

INVOKAMET is not recommended for use in patients under 18 years of age.

Adults aged 65 years of age and older

You could have more side effects with INVOKAMET.

Check-ups and testing

You will have regular visits with your healthcare professional before and during treatment with INVOKAMET to monitor your health. They may check:

- your blood sugar levels. INVOKAMET will cause your urine to test positive for sugar.
- that your kidneys are working properly.
- blood fat levels.
- the amount of red blood cells in your blood; and
- the potassium levels in your blood.
- ketone levels in your blood or urine. Ketones are a type of chemical that your liver produces when it breaks down fats for energy.
- the levels of lithium in your blood.
- your blood levels of vitamin B₁₂.

Broken bones (fracture)

Taking INVOKAMET may increase your risk of breaking a bone. Talk to your doctor about factors that increase the risk for broken bones.

Surgery and illnesses

Tell your healthcare professional if you:

- have a serious medical illness.
- are going to have a surgery.
- are hospitalized for a serious infection or a serious medical illness; or
- had major surgery.

Stopping treatment

- Your healthcare professional may temporarily stop your INVOKAMET treatment:
 - before and after certain types of surgery, or procedures associated with long periods of fasting. Your healthcare professional may stop your treatment for at least three days before you undergo these procedures or surgeries. or
 - when you are hospitalized for a serious infection or illness

If INVOKAMET is stopped, your healthcare professional will:

- continue to monitor for signs or symptoms of DKA.
- tell you when to start taking INVOKAMET again.

Driving and using machinery

INVOKAMET may cause dizziness or light-headedness. Do NOT drive or use machines until you know how the medicine affects you.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines

The following may interact with INVOKAMET:

- digoxin, used to treat heart problems.
- furosemide or other diuretics (water pills), used to treat high blood pressure and other heart problems.
- insulin or a sulfonylurea (such as glimepiride, gliclazide, or glyburide), used to help control blood sugar.
- carbamazepine or phenobarbital, used to treat seizures.
- barbiturates, used as sedatives and sleep-aids.
- efavirenz or ritonavir, used to treat HIV infection.
- rifampin, an antibiotic used to treat bacterial infections such as tuberculosis.
- lithium, used to treat bipolar disorder.
- St. John's wort, an herbal product used to treat depression.
- Angiotensin-Converting Enzyme (ACE) inhibitors, Angiotensin Receptor Blockers (ARB) used to treat high blood pressure.
- phenprocoumon and other drugs used prevent blood clots and thin the blood.
- other medicines that can cause high blood sugar, and may lead to a loss of blood sugar control, such as:
 - corticosteroids such as prednisone, used to treat a broad range of diseases
 - isoniazid, used to treat tuberculosis

- phenothiazines, a group of medicines used to treat mental problems, including schizophrenia
- thiazide (water pills), used to treat high blood pressure
- thyroid medicines such as levothyroxine
- birth control pills and other products containing estrogens
- a group of medicines known as calcium channel blockers, used to treat heart problems such as: nifedipine, amlodipine, felodipine, verapamil, diltiazem
- medicine used to treat epilepsy (seizures) such as phenytoin
- nicotinic acid, used to prevent and treat niacin deficiency
- bronchodilators (known as beta-2-agonists) medicines that make breathing easier, used to treat asthma like salbutamol or formoterol
- alcohol.

How to take INVOKAMET:

- Take INVOKAMET exactly as your healthcare professional tells you. Check with your healthcare professional if you are not sure.
- Take twice per day with food. This is to reduce your chance of having an upset stomach.
- Swallow the tablet(s) whole.
- Continue taking INVOKAMET for as long as your healthcare professional tells you to.
- Your healthcare professional may temporarily stop your treatment for a minimum of three days if you are undergoing:
 - a major surgery.
 - procedures that are associated with long periods of fasting.

Your healthcare professional may prescribe INVOKAMET alone or together with another medicine to help control your blood sugar.

Usual dose:

Your healthcare professional will decide the best dose that is right for you.

Overdose:

If you think you, or a person you are caring for, have taken too much INVOKAMET, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed dose:

- If you forget to take a dose of INVOKAMET, take it as soon as you remember. However, if it is nearly time for the next dose, skip the missed dose.
- Do not take a double dose to make up for a forgotten dose.

What are possible side effects from using INVOKAMET?

These are not all the possible side effects you may feel when taking INVOKAMET. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- Changes in taste or a metallic taste
- Changes in urination:
 - urinating more often or in larger amounts
 - an urgent need to urinate
 - a need to urinate at night
- Constipation, excess gas, abdominal discomfort
- Fatigue
- Feeling thirsty
- Headache
- Low vitamin B₁₂ (vitamin B₁₂ deficiency)
- Nausea, vomiting, diarrhea, indigestion, loss of appetite
- Rash, hives

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Vaginal yeast infection: vaginal odor, white or yellowish vaginal discharge, and/or itching		✓	
COMMON			
Balanitis (yeast infection of the penis): rash or redness of the penis or foreskin		✓	
Bone fracture (broken bone)		✓	
Skin ulcer (a break or sore on the skin with tissue breakdown) predominantly of the lower legs: It may start off red then get swollen and tender. Next, blisters can form with loss of skin layers. It can lead to an open round crater with a bad smell. Ulcers take a long time or may not heal.		✓	
Urinary tract infection: burning sensation when urinating, cloudy or bloody urine, strong odor		✓	
UNCOMMON			
Dehydration (not having enough water in your body): dry or sticky mouth, headache, dizziness or urinating less often than normal		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Hypotension (low blood pressure): fainting, dizziness or light-headedness with standing		✓	
Kidney problems: any change in the amount, frequency or colour (pale or dark) of urine		✓	
Peripheral ischemia (blocked or narrow blood vessels): Leg pain with walking that gets better with rest. Poor circulation, bluish, cold skin, and poor nail and hair growth. It can lead to Skin Ulcers and Lower Leg or Toe Amputation .		✓	
RARE			
Acute kidney infection: painful, urgent or frequent urination, lower back (flank) pain, fever or chills, cloudy or foul smelling urine, blood in your urine			✓
Angioedema (swelling in deep layers of skin) and anaphylactic reaction (severe allergic reactions): rash, hives, swelling of the face, eyes, lips or throat, difficulty swallowing or breathing, wheezing, fever, stomach cramps, chest discomfort or tightness, unconsciousness			✓
Diabetic ketoacidosis (when your body produces high levels of blood acids called ketones): difficulty breathing, feeling very thirsty, vomiting, stomach pain, nausea, loss of appetite, confusion, and unusual tiredness, a sweet smell to the breath, a sweet or metallic taste in the mouth or a different odour to urine or sweat			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Encephalopathy (disease of the brain that severely alters thinking): Possible neurological symptoms include: muscle weakness in one area, poor decision-making or concentration, involuntary twitching, trembling, difficulty speaking or swallowing, seizures.			✓
Fournier's gangrene (a serious infection affecting soft tissue around the groin): pain or tenderness, redness of the skin, or swelling in the genital or perineal area, with or without fever or feeling very weak, tired, or uncomfortable			✓
Hemolytic anemia (breakdown of red blood cells): symptoms may include fatigue, pale color, rapid heartbeat, shortness of breath, dark urine, chills, and backache.			✓
Severe hypoglycemia (severe low blood sugar): disorientation, loss of consciousness, seizure			✓
Lactic acidosis (a build-up of lactic acid in your blood): feeling cold or uncomfortable, severe nausea with or without vomiting, stomach pain, unexplained weight loss, rapid breathing			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Peripheral neuropathy (a result of damage to your peripheral nerves): signs and symptoms might include gradual onset of numbness, prickling or tingling in your feet or hands, which can spread upward into your legs and arms, sharp, jabbing, throbbing, freezing or burning pain, extreme sensitivity to touch, lack of coordination and falling, muscle weakness or paralysis if motor nerves are affected.			✓
Urosepsis (severe infection that spreads from the urinary tract and throughout body): fever or low body temperature, rapid breathing, chills, rapid heartbeat, pain with urination, difficulty urinating, frequent urination			✓
VERY RARE			
Liver problems: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		✓	
Pancreatitis (inflammation of the pancreas): severe stomach pain that lasts and gets worse when you lie down, nausea, vomiting		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store INVOKAMET in the original container.
- Store at room temperature (15-30°C).
- Keep out of reach and sight of children.

Do not use INVOKAMET after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

If you want more information about INVOKAMET:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website www.janssen.com/canada or call 1-800-567-3331 or 1-800-387-8781.

This leaflet was prepared by: JANSSEN Inc., Toronto, Ontario, M3C 1L9

Last Revised: June 13, 2024

All trademarks used under license.